

# Acute Phase Response in Animals: A Review

Carolyn Cray,<sup>1\*</sup> Julia Zaias,<sup>1,2</sup> and Norman H Altman<sup>1</sup>

The acute phase response is a complex systemic early-defense system activated by trauma, infection, stress, neoplasia, and inflammation. Although nonspecific, it serves as a core of the innate immune response involving physical and molecular barriers and responses that serve to prevent infection, clear potential pathogens, initiate inflammatory processes, and contribute to resolution and the healing process. Acute phase proteins, an integral part of the acute phase response, have been a focus of many applications in human diagnostic medicine and recently have been identified in common animal species. Potential applications to diagnosis, prognosis, assessment of animal health, and laboratory animal welfare are readily apparent.

**Abbreviations:** APP, acute phase protein; APR, acute phase response; CRP, C reactive protein; SAA, serum amyloid A.

Acute phase proteins (APP) are blood proteins primarily synthesized by hepatocytes as part of the acute phase response (APR). The APR is part of the early-defense or innate immune system, which is triggered by different stimuli including trauma, infection, stress, neoplasia, and inflammation. The APR results in a complex systemic reaction with the goal of reestablishing homeostasis and promoting healing. The APR has been referred to as the 'molecular thermometer' whereby quantitation of individual APP can provide an assessment of the response to the triggering event.<sup>10,82,93</sup>

APP have been well recognized for their application to human diagnostic medicine and have been described to have value in the diagnosis and prognosis of cardiovascular disease, autoimmunity, organ transplant, and cancer treatment.<sup>15,16,24,36,50,57,73,102</sup> C-reactive protein (CRP) was the first APP described (in the early 1930s), and its presence early during pneumococcal infection of monkeys and humans thereby led to the coinage of 'acute phase' term.<sup>1,123</sup> Today, CRP remains an APP of primary interest in humans, where it is a major marker of infection, autoimmune disease, trauma, malignancy, and necrosis including myocardial infarction.<sup>92</sup> Furthermore, CRP has been proposed as a marker for wellness assessments, which is a common role proposed in many studies of human and animal APP.<sup>73</sup>

Although the study of serum proteins in animals was commonly conducted in research investigations in the mid-1900s, the wider application of APP in veterinary medicine was not reported until the early 1990s. Several review papers with descriptions of APP and APR in companion animals and rodents are available.<sup>10,39,82,93</sup> Like human APP, animal APP have been well documented to be sensitive to similar triggering events. In large animals, APP have been further proposed as markers of 'herd health'.<sup>40,82,93</sup> In companion animals, APP have especially been identified for prognostic applications.<sup>10</sup> The use of APP in laboratory animals has essentially been limited to research based investigations, but APP have been proposed as an objective marker

for humane experimental endpoints.<sup>39,89</sup> The following sections summarize the biologic effects of the APR, current issues and options in the quantitation of APP and APR, applications of APP in the diagnosis and prognosis of disease in animal species, and proposed applications of APP in laboratory animals.

## The Acute Phase Response


Innate immunity fulfills an important role in the body's early defense mechanism and serves to initiate the acquired immune response. The innate immune system is very diverse and includes physical barriers, phagocytes, complement, and toll-like receptors which serve to prevent infection, eliminate potential pathogens, and initiate the inflammatory process.<sup>64</sup> Later aspects of induced innate responses are dependent on cytokines and chemokines, which are generated by activated cells including monocytes, macrophages, fibroblasts, endothelium, platelets, keratinocytes, and T cells.<sup>64</sup> These proinflammatory signals, including IL1, TNF $\alpha$ , and IL6, have numerous effects throughout the body including inducing the APR (Figure 1). The APR is a core part of the innate immune response and is observed across all animal species. In fact, counterparts of mammalian APP have been identified in invertebrates and fish, in which the APR is proposed to be more robust than that in mammals to compensate for a less-evolved adaptive immune response.<sup>4,100</sup>

The APR may result in changes in more than 200 proteins grouped as either positive APP or negative APP.<sup>72</sup> In nearly all animal species, albumin represents the major negative APP which, during the APR, decreases in blood concentration and may represent either selective loss of albumin due to renal or gastrointestinal changes or a decrease in hepatic synthesis.<sup>72</sup> The decrease in albumin synthesis is postulated to allow for the unused pool of amino acids to instead be used to generate positive APP and other important mediators of inflammation.<sup>90</sup> Positive APP are those that increase during the APR. They are further classified as major, moderate, or minor, depending on the magnitude of increase during the APR. Traditionally, major proteins represent those that increase 10- to 100-fold, moderate proteins represent those that increase 2- to 10-fold, and minor proteins represent those with only a slight increase.<sup>10</sup> These classifications may vary

Received: 27 Jul 2009. Revision requested: 23 Aug 2009. Accepted: 05 Sep 2009.

<sup>1</sup>Division of Comparative Pathology and <sup>2</sup>Division of Veterinary Resources, University of Miami Miller School of Medicine, Miami, Florida.

\*Corresponding author. Email: c.cray@miami.edu



**Figure 1.** The acute phase response. The figure reflects information drawn from references 10, 82, and 93.

slightly within different APR publications, although the implied changes by category remain the same.<sup>90</sup> Major proteins often are observed to increase markedly within the first 48 h after the triggering event and often have a rapid decline due their very short half-life.<sup>70</sup> Moderate and minor proteins follow in magnitude of response and may both increase more slowly and be more prolonged in duration, depending on the status of the triggering event. Moderate and minor APP may be observed more often during chronic inflammatory processes.<sup>10,93</sup> Species differences are readily apparent in the classification of positive APP (Figure 2).

The biologic functions of APP are vast and have been reviewed.<sup>10,70,82,93,116</sup> A summary of these key observations follows. For example, CRP can act as an opsonin by binding residues and polysaccharides on bacteria, fungi, and parasites to activate complement and phagocytosis. In addition, CRP can both up-regulate and downregulate cytokine production and chemotaxis. Serum amyloid P is thought to be an analog to CRP in some species. Serum amyloid A (SAA), another APP often classified as a major protein, has been demonstrated to result in chemotaxis of monocytes, polymorphonuclear cells, and T cells. In addition, SAA has marked inhibitory effects and is assumed to be important in the down regulation of the inflammatory process. Haptoglobin reduces oxidative damage associated with hemolysis by binding free hemoglobin. In addition, it has been observed to have bacteriostatic and immunomodulatory effects. The APP  $\alpha$ 1-acid glycoprotein binds LPS and inhibits its activity;  $\alpha$ 1-acid glycoprotein also can bind to numerous drugs and may sustain drug transport levels even with a decrease in albumin, the negative acute phase protein. Whereas  $\alpha$ 1-acid glycoprotein can also downregulate neutrophils and complement,  $\alpha$ 2-macroglobulin is 1 of several APP that have protease inhibitory activity and are important in removing enzymes released during injury. Major acute phase protein (porcine) (or pig-MAP) has been identified

in porcine species and inhibits trypsin. Ceruloplasmin scavenges free radicals. Fibrinogen provides a substrate for fibrin formation and thus is important in tissue repair. Transferrin is proposed to be a positive APP in avian species, although in most mammals, it is considered a negative APP. In summary, positive APP all have multiple functions including modulating the immune system, protein transport, and tissue protection from damage by the inflammatory process.

### Assays of Acute Phase Proteins

In basic health assessments, total protein and albumin can be measured on is commonly measured on automated chemistry analyzers, and globulin is a value calculated from these measurements. In human and veterinary medicine, protein electrophoresis is a diagnostic tool that has widely been applied to the study of the APR for more than 40 y.<sup>2,17,72,107</sup> Although rarely solely diagnostic of a particular disease, protein electrophoresis is an excellent method for the detection of acute and chronic inflammatory processes and stimulation of humoral immunity. Veterinary applications include its ancillary use in the diagnosis of feline infectious peritonitis, ehrlichiosis, and myeloma and, more recently, in the diagnosis and prognosis of infectious diseases in avian species.<sup>19,20,72</sup>

The general technical principle of protein electrophoresis involves overlaying plasma or serum on a thin agarose gel. An electric current is applied to the gel, which causes the proteins to migrate according to their charge and size. The movement of the proteins creates bands in the gel, which can be quantitated by densitometry. Protein electrophoresis can be performed by using commercially available systems, which allow for on-demand performance, are relatively inexpensive, require only a few microliters of serum, and provide results within a few hours. In addition, multiple samples can be analyzed simultaneously, and overall changes in the APR can be quantitated. The proteins that are resolved by protein electrophoresis are albumin and the 4 fractions of globulins referred to as  $\alpha$ 1,  $\alpha$ 2,  $\beta$ , and  $\gamma$ . Each globulin fraction is composed of APP or immunoglobulins (or both). Protein electrophoresis provides a more accurate reflection of the albumin:globulin ratio, given that as albumin methodologies in chemistry analyzers are optimized for human albumin and that bromocresol green can bind animal globulins with extended reaction times.<sup>41,115</sup> Therefore, protein electrophoresis can provide more accurate albumin quantitation and visualization of globulin fractions to monitor the overall progression of the APR. Protein electrophoresis does not enable quantitation of single proteins but rather of groups of proteins that are mediators of acute inflammatory processes. More than 200 blood proteins have been defined and are likely present in protein electrophoresis fractions.<sup>72</sup> Changes in proteins with low concentrations may not be detected by this technique. Protein electrophoresis electrophoretograms from commercial systems are thought to reveal changes in approximately 30 major serum proteins.  $\alpha$ 1 globulins include  $\alpha$ 1-antitrypsin and  $\alpha$ 1-acid glycoprotein,  $\alpha$ 2 globulins include  $\alpha$ 2-macroglobulin and haptoglobin;  $\beta$  globulins include transferrin, SAA, fibrinogen, and CRP; and  $\gamma$  globulins primarily comprise IgG.<sup>72</sup> Although reference intervals are available for most common mammalian species, these ranges often are derived from nonlaboratory animals and thus should be evaluated critically before their application. We recently published normal reference intervals for common inbred strains of mice, including age-relative

Species	Major (>10-fold increase)	Moderate (1- to 10-fold increase)
Cat	$\alpha$ 1-acid glycoprotein, serum amyloid A	haptoglobin
Chicken	none	$\alpha$ 1-acid glycoprotein, ceruloplasmin, serum amyloid A, transferrin
Cow	haptoglobin, serum amyloid A	$\alpha$ 1-acid glycoprotein, C-reactive protein, fibrinogen
Dog	C-reactive protein, serum amyloid A	$\alpha$ 1-acid glycoprotein, ceruloplasmin, haptoglobin
Goat	haptoglobin, serum amyloid A	fibrinogen
Horse	serum amyloid A	fibrinogen, haptoglobin
Human	C-reactive protein, serum amyloid A	$\alpha$ 1-acid glycoprotein, fibrinogen, haptoglobin
Mouse	haptoglobin, serum amyloid A, serum amyloid P	C-reactive protein, fibrinogen
Nonhuman Primates	C-reactive protein	$\alpha$ 2-macroglobulin, fibrinogen, serum amyloid A
Pig	haptoglobin, serum amyloid A, major acute phase protein	$\alpha$ 1-acid glycoprotein
Rabbit	haptoglobin, serum amyloid A	$\alpha$ 1-acid glycoprotein, C-reactive protein, fibrinogen
Rat	$\alpha$ 1-acid glycoprotein, $\alpha$ 2-macroglobulin	C-reactive protein, fibrinogen, haptoglobin
Sheep	haptoglobin, serum amyloid A	$\alpha$ 1-acid glycoprotein, C-reactive protein

**Figure 2.** Major and moderate acute phase proteins in different animal species. The figure reflects information drawn from references 10, 13, 39, 59, 82, 93, and 108.

reference intervals, by using a commercially available agarose-based electrophoresis system.<sup>137</sup> Investigators planning to implement protein electrophoresis or APP as tools in laboratory animal medicine should consider establishing a baseline for species and strains, given that the variation present in normal resting values likely is unknown.

Individual APP have been documented in many species by using ELISA, radioimmunoassay, nephelometry, and immunoturbidimetry; Western blot; and mRNA analyses.<sup>10,29,39,90,108</sup> Unfortunately, due to wide species variation in APP, few cross-reactive reagents are available. Assays for human APP have variable cross-reactivity.<sup>31,38</sup> Ideally, species-specific assays should be used. Although some commercially available ELISA kits are available to measure specific APP including CRP, SAA, and haptoglobin, most of these assays lack automation and are expensive. The ELISA methodology is best suited for batch analysis of many samples to make best use of technical labor and reagents. Some automated assays marketed for human APP have been validated for animal use, and few have been implemented for species-specific veterinary use.<sup>29,60,61,122</sup> Serum amyloid A appears to be conserved within mammalian species, and haptoglobin can be quantitated based on its universal reactivity with hemoglobin.<sup>52,61,122</sup> Standardization of APP assays has been a focus of international veterinary meetings and publications.<sup>32,111</sup> Species-specific controls should be considered when possible to allow for standardization of such assays between laboratories.<sup>32,111</sup> Other technical issues with APP testing include the variations in APP concentrations observed with different anticoagulants and interfering sample conditions such as hemolysis and lipemia.<sup>10</sup> Although APP are reported to be stable at  $-20^{\circ}\text{C}$ , for prolonged storage,  $-70^{\circ}\text{C}$  is recommended.<sup>10</sup>

Few studies in animals directly compare protein electrophoresis and APP ELISA assays in their ability to gauge the APR. In a study following rats for 21 d after dosing with complete Fre-

und adjuvant, haptoglobin increased within 12 h.<sup>43</sup> This increase, nearly 700% of normal values, occurred concurrently with a significant decrease in albumin and increases in  $\alpha$ 2 and  $\beta$  globulins as quantitated by protein electrophoresis. Alterations in globulin, albumin, and haptoglobin concentration continued to be present until day 21, when normal levels were observed in all parameters. Therefore, although they are different methodologies, protein electrophoresis and APP ELISA could provide views of APR that reflect the same time course. The relative sensitivity of these methods may be better gauged by using less aggressive inflammatory stimuli or a more diverse inflammatory trigger, such as a chronically progressive disease condition.

### Applications of Acute Phase Proteins in Diagnosis and Prognosis

Before the advent of specific APP assays, monitoring the albumin:globulin ratio had been a standard in human and veterinary medicine to monitor inflammatory processes. Technology has expanded from protein electrophoresis to species-specific assays, which offer a more detailed examination of APP and their changes with disease progression. Triggering events of inflammation, infection, trauma, neoplasia, and stress alter APP levels in animal species; an extensive review of these publications is presented in Figure 3, which highlights APP changes in both experimental conditions and naturally occurring disease. For most species, traditional inflammatory agents such as croton oil and turpentine have been used to assess the types of APP produced and their magnitude and kinetics of expression. Changes in APP in experimental models of infection and naturally occurring inflammatory and infectious disease also have been assessed. Furthermore, several studies have demonstrated the utility of APP quantitation in monitoring neoplasia and stress.

Triggering event	Animal	Significant changes in acute phase proteins	References
Infection	Cat	Serum amyloid A increased with feline infectious peritonitis infection	44, 118
		$\alpha$ 1-acid glycoprotein and haptoglobin increased with feline infectious peritonitis or feline immunodeficiency virus infection	28
	Chicken	Transferrin increased with <i>Staphylococcus aureus</i> infection	12
		$\alpha$ 1-acid glycoprotein increased with experimental infection with infectious bursal disease virus	58
	Cow	Haptoglobin, serum amyloid A, and fibrinogen increased with infections including bovine coronavirus and bovine adenovirus	3, 40
	Dog	C-reactive protein and serum amyloid A levels correlate with the severity of pyometra	21, 85
		C-reactive protein and ceruloplasmin increased with babesiosis	127
		C-reactive protein increased in dogs experimental infected with <i>Ehrlichia canis</i>	110
		C-reactive protein increased in dogs naturally infected with <i>Leishmania</i> spp.; higher in symptomatic dogs	79
	Horse	Serum amyloid A increased with colic	130
		Serum amyloid A increased with bacterial infection	51, 55
	Mouse	Serum amyloid P and haptoglobin increased experimental infection with <i>Trypanosoma</i>	33
		Serum amyloid P increased in experimental malaria infection	6
		Serum amyloid A and serum amyloid P increased with bacterial pneumonia	99
	Nonhuman Primate	C-reactive protein increased in macaques inoculated with <i>Bordatella bronchispetica</i>	66
	Ovine	C-reactive protein, haptoglobin, ceruloplasmin, and fibrinogen increased with bacterial infection	112, 128
Ceruloplasmin, fibrinogen, and haptoglobin increased and albumin decreased with yeast infection		95	
Haptoglobin, serum amyloid A, and $\alpha$ 1-acid glycoprotein increased in an experimental model of bacterial lymphadenitis		34	
Pig	C-reactive protein and serum amyloid A increased with experimental infection with the respiratory pathogen <i>Actinobacillus pleuropneumoniae</i>	113	
	Major acute phase protein and haptoglobin increased with pig coronavirus infection and wasting syndrome	46	
	C-reactive protein, serum amyloid A, and haptoglobin are valuable markers in natural infection with different porcine viruses and <i>Mycoplasma</i> infection	91	
Inflammation	Cat	Serum amyloid A increased in acute pancreatitis in cats	118, 119
		Serum amyloid A, $\alpha$ 1-acid glycoprotein, and haptoglobin increased after surgery or injection with either LPS or turpentine	71
	Chicken	Transferrin increased with injection of croton oil or turpentine	12, 124, 132
		$\alpha$ 1-acid glycoprotein increased with injection of <i>Escherichia coli</i> LPS	84, 117
	Dog	C-reactive protein and $\alpha$ 1-acid glycoprotein increased with injection of turpentine oil; 14-d time course for recovery	48, 136
		Haptoglobin and fibrinogen significantly increased with hyperadrenocorticism	8
		C-reactive protein correlated to severity and prognosis of acute pancreatitis	78, 85
		C-reactive protein levels correlated with inflammatory bowel disease activity	65
		C-reactive protein, fibrinogen, haptoglobin, and serum amyloid A increased with experimentally induced gastric mucosal injury	7
		C-reactive protein increased after surgery; levels decrease by time of suture removal	133
		C-reactive protein increased with chronic valvular disease	103
		C-reactive protein increased with canine idiopathic polyarthritis; good prognostic indicator	85, 88
		C-reactive protein increased with panniculitis	85
	Horse	Serum amyloid A increased and correlative to severity of joint disease	63
		Serum amyloid A increased after surgery	97
		Serum amyloid A increased after experimentally induced noninfectious arthritis	56, 62

Figure 3. Summary of acute phase responses by triggering event.

The dichotomy of both rapid- and slow-responding APP provides an added dimension in clinical interpretation that is not readily seen with changes in other blood analytes and complete blood counts. In a large study, clinically ill cattle were assessed

on the basis of signs, diagnostic testing, and postmortem findings.<sup>53</sup> The animals were categorized as experiencing either acute or chronic inflammatory processes. Serum amyloid A levels had 100% sensitivity and haptoglobin had 76% specificity in distin-

Triggering event	Animal	Significant changes in acute phase proteins	References
	Mouse	Serum amyloid A increased with injection of ethanol	98
		Serum amyloid A increased with injection of croton oil or <i>Aspergillus</i> antigen	74
		Serum amyloid A, serum amyloid P, and $\alpha$ 1-acid glycoprotein increased with injection of hepatotoxic agent or LPS	45, 75, 131
		Serum amyloid A and serum amyloid P increased in IL2 <sup>-/-</sup> mice which develop severe colonic disease	22
	Nonhuman Primate	Serum amyloid A increased in macaques with subclinical secondary hepatic amyloidosis	54, 77
		Fibrinogen and $\alpha$ 2-macroglobulin increased in rhesus monkeys treated with IL6	83
		C-reactive protein increased in different monkey species injected with turpentine oil	67
	Pig	C-reactive protein and haptoglobin increased with injection of turpentine	35
		C-reactive protein, serum amyloid A, and haptoglobin increased in pigs with tail and ear bites and arthritis	91
	Rabbit	Serum amyloid A increased in an experimental model of non infectious chronic inflammation	101
		C-reactive protein increased with injection of turpentine	42
	Rat	$\alpha$ 1-acid glycoprotein, $\alpha$ 2-macroglobulin, fibrinogen, and haptoglobin increased and albumin decreased with injection of turpentine	9, 69, 108
		$\alpha$ 2-macroglobulin increased 25-fold in rats subjected to surgery	68
$\alpha$ 1-acid glycoprotein increased with injection of prostaglandin E2 for induction of fever		129	
Haptoglobin increased with injection of complete Freund adjuvant or carbon tetrachloride		43	
Haptoglobin increased and albumin decreased with injection of phosphodiesterase inhibitor		27	
Sheep	Ceruloplasmin and fibrinogen increased in animals with artificial pulmonary obstruction	94	
Neoplasia	Cat	Serum amyloid A increased with lymphoma and malignant mesothelioma	118
		$\alpha$ 1-acid glycoprotein increased with lymphoma	18
		$\alpha$ 1-acid glycoprotein increased with carcinoma, sarcoma, and round cell tumors	109
	Dog	C-reactive protein useful in determining remission status in cases of multicentric lymphoma	87
		C-reactive protein and haptoglobin increased with lymphatic neoplasia	80
		C-reactive protein increased with hemangiosarcoma	85
	C-reactive protein increased with mammary tumors; albumin decreased in metastasis	120	
	C-reactive protein, haptoglobin, and ceruloplasmin increased with hematologic and neoplastic diseases	121	
Mouse	C-reactive protein and haptoglobin increased in transgenic mice with intestinal adenoma	37	
	Serum amyloid A and C-reactive protein increased in model of fibrosarcoma	14	
Stress	Cow	serum amyloid A increased in cows living in semiferal conditions with nutritional stress	104
		serum amyloid A increased after exposure to complex stress (transportation, tie stall housing, slippery floors, social isolation)	76
		serum amyloid A increased in calves after 3-h truck transportation	5
		fibrinogen increased in calves with abrupt weaning	49
	Pig	C-reactive protein and haptoglobin increased after 48-h truck transportation; major acute phase protein increased after 24 h in both poor and superior conditions of truck transportation	96, 106
		major acute phase protein increased after tail biting	106
		C-reactive protein, haptoglobin, and major acute phase protein increased after repeated-bleeding protocols (sensitization to procedure)	106
	Rat	$\alpha$ 1-acid glycoprotein and haptoglobin increased after exposure to tail shock	23

Figure 3. Continued

guishing the groups. In comparison, neutrophil counts had much lower sensitivity and specificity (30% to 70%). In sheep, increases in haptoglobin had 85% sensitivity and specificity compared with 52% sensitivity and 75% specificity of WBC counts in those animals with naturally occurring bacterial infection.<sup>112</sup> In a retrospective evaluation of inflammatory conditions in dogs, APP showed significant changes in the absence of changes in the total or differential WBC count.<sup>114</sup> In response to steroid treatment, dogs with

idiopathic polyarthritis showed a positive response to treatment (decreased symptoms) as well as significant decreases in CRP whereas total WBC count remained unchanged.<sup>88</sup> In a review of more than 900 cases of inflammation in dogs with various diseases, CRP concentrations were significantly correlated with disease, whereas only slight or no correlation was found with total WBC and band neutrophil counts.<sup>85</sup> In a cat with pancreatitis, SAA was increased with onset of symptoms whereas the WBC count was

normal.<sup>119</sup> With rapid resolution of the symptoms in this cat, SAA returned to normal levels, whereas the WBC count was just beginning to increase. In total, these findings are consistent with the expected delay in marked WBC changes due to the need to generate cells in the bone marrow, whereas APP concentrations can increase within hours.

Several studies have shown an association between APP levels and the severity of disease. In a study of naturally occurring babesiosis, CRP levels were significantly higher in dogs with severe or complicated disease.<sup>127</sup> In canine leishmaniasis, CRP was increased in symptomatic but not asymptomatic animals.<sup>79</sup> One group of investigators proposed a canine inflammatory bowel disease activity index, which was based on a panel of clinical symptoms relative to disease activity in canine inflammatory bowel disease.<sup>65</sup> Increased levels of CRP and  $\alpha$ 1-acid glycoprotein were associated with increasing activity indices of disease.

In these aforementioned studies as well as many others, APP have been demonstrated to change with a positive prognosis. Of note, the half-life of most APP is only 24 to 48 h. Thus, changes in APP may be more sensitive indicators of healing and resolution than other clinical tests. After surgery in dogs, CRP levels were found to decline and mirror the disappearance of clinical signs when the total WBC count continued to be elevated.<sup>21,85</sup> As mentioned earlier, CRP levels were decreased with a positive response to steroid treatment in dogs with polyarthritis.<sup>88</sup> In addition, after chemotherapy for multicentric lymphoma in dogs, CRP was within normal reference intervals in cases with complete remission.<sup>87</sup> In the study of canine inflammatory bowel disease, levels of CRP decreased with a positive response to therapy.<sup>65</sup> Haptoglobin, CRP, and SAA were monitored in dogs with pyometra before and after surgical sterilization.<sup>21</sup> Compared with those of normal dogs before surgery, CRP and SAA values in dogs with pyometra peaked within 24 h after surgery and then declined slowly in dogs that recovered without complications.

Several authors have suggested that a single APP should not be used exclusively to monitor a disease process. Instead, an APP index has been used in both human and veterinary medicine.<sup>47</sup> This index includes both positive and negative APP, as well as APP that increase both rapidly and slowly, thereby forming a comprehensive index that would correlate with the severity of the inflammatory process. A recent editorial on APP interpretation in companion animals summarized several integral points, including the need for an APP profile (for example, multiple positive APP).<sup>11,126</sup>

## Potential Novel Applications of Acute Phase Proteins in Laboratory Animal Medicine

Defining humane endpoints to experimental protocols remains a difficult task. Refinement of animal protocols is a primary goal of both investigators and institutional animal care and use committees, not only to gain valuable experimental information from animals prior to morbidity and mortality but also to meet the important obligation of minimizing pain and stress. Definition of humane endpoints often is driven by subjective measures of clinical condition, frequently in combination with weight loss. In many animal protocols, a moribund condition is used as a point of experiment termination. In some cases, a scoring of clinical and behavioral signs can be used to indicate that euthanasia is warranted. Such signs can include ruffling of the hair or coat,

hunched posture, lethargy, decreased consumption of food or water, weight loss, diarrhea, bleeding, respiratory difficulty, impaired mobility, and unconsciousness.<sup>86,125</sup> In many cases, such clinical signs do not appear in all of the experimental animals, and some of these clinical signs may be transient and precede eventual recovery of the animal.<sup>86</sup> Another prevailing problem is that many of the variables monitored are subjective, and the accuracy of these assessments can vary among laboratory personnel and veterinary staff members. Moreover, investigators working with new models or newly created transgenic mice may not be able to rely on scoring systems determined previously on other models.<sup>25,26</sup> With the increasing use of animals in research, including the advent of transgenic mice and the increased need for models for translational studies, guidelines that are more objective are essential. APP have been proposed to be biochemical markers of stress, infection, and pain in laboratory animals.<sup>89</sup> Although no APP study has assessed this particular application specifically, a role for quantitating APP can be envisioned, given the broad increases that follow with triggering events including infection, inflammation, trauma, stress, and neoplasia, many of which are common conditions in experimental protocols involving the use of laboratory animals.

APP have been used to gauge stress with regard to animal well being. One investigator proposed a link among catecholamines, glucocorticoids, and production of APP in animals.<sup>81</sup> APP can indicate stress in pigs and calves. Pigs shipped under typical conditions had higher major acute phase protein (porcine), haptoglobin, SAA, and CRP than did pigs that were shipped under conditions that included the provision of sawdust, water, and feed.<sup>96</sup> Similar studies also demonstrated changes in APP in pigs, newly weaned calves, and cattle exposed to with new housing situations,<sup>5,49,76,106</sup> and in pigs that were unaccustomed to various handling procedures.<sup>106</sup> These results indicate the possible use of APP to monitor stress in laboratory animals and thus provide another marker of animal well being. Stressors that could be assessed in future studies of laboratory animals include physical environment (lighting, temperature, and noise), housing type, husbandry and handling, and shipping.

Another potential use of APP in laboratory animals is assessment of the general health of a colony. As indicated in Figure 3, APP are sensitive markers of inflammation and infection in animal species. In large animals, APP have been proposed as markers of herd health, through which a large group of animals may be assessed for food safety and herd management.<sup>40,126</sup> In 1 study, 2 groups of calves were examined over an extended time period. The group that had a higher incidence of disease also demonstrated significantly higher levels of APP.<sup>40</sup> Dogs maintained as laboratory animals had significantly lower CRP levels than did those kept as companion animals.<sup>134,135</sup> The authors hypothesized that the husbandry conditions and isolation of the laboratory animals minimized contact with potential inflammatory triggers.<sup>10</sup>

The use of APP to monitor rodent colony health may have limited value. Monitoring of APP levels would not be expected to be a primary tool used preferentially over traditional serologic and PCR screening methods; rather, APP assessment potentially would be an ancillary test. Additional studies should be performed before the implementation of APP monitoring in rodent colonies. For example, APP may be more suited to monitoring the health of individual animals. We have previously reported increased levels of  $\alpha$  and  $\beta$  globulins in transgenic mice with der-

matitis.<sup>137</sup> In addition, APP might be used to screen animals (cat, dog, nonhuman primate, pig, and rabbit) during quarantine or before entry into experimental protocols. The nonspecificity of the APR has been proposed to offer a possible 'snapshot' of animal health and thus may reflect the presence of subclinical disease or other management issues.<sup>30,105</sup> APP may help to further characterize similar issues in laboratory animal colonies.

## Conclusions

The advantages and uses of APP assays are well supported in the human and veterinary medical literature. Unfortunately, because of the practical limitations of current technology, clinical application of APP analysis is not widespread. Continuing challenges include the need for automated assays and standardization of tests across laboratories. However, many potential uses are possible for APP and APR in laboratory animal medicine. Exceptionally sensitive yet advantageously nonspecific markers of diverse inflammatory etiologies, APP are excellent candidates through which to better study animal models of disease, monitor animal health, and objectively assess animal wellbeing.

## References

- Abernethy JT, Avery OT. 1941. The occurrence during acute infections of a protein not normally present in the blood. I. Distribution of the reactive protein in patients' sera and the effect of calcium on the flocculation reaction with C polysaccharide of *Pneumococcus*. *J Exp Med* 73:173–182.
- Alper CA. 1974. Plasma protein measurements as a diagnostic aid. *N Engl J Med* 291:287–290.
- Angen O, Thomsen J, Larsen LE, Larsen J, Kokotovic B, Heegaard PM, Enemark JM. 2009. Respiratory disease in calves: microbiological investigations on transtracheally aspirated bronchoalveolar fluid and acute phase protein response. *Vet Microbiol* 137:165–171.
- Armstrong PB, Quigley JP. 1999. Alpha2-macroglobulin: an evolutionarily conserved arm of the innate immune system. *Dev Comp Immunol* 23:375–390.
- Arthington JD, Eichert SD, Kunkle WE, Martin FG. 2003. Effect of transportation and commingling on the acute-phase protein response, growth, and feed intake of newly weaned beef calves. *J Anim Sci* 81:1120–1125.
- Balmer P, McMonagle F, Alexander J, Phillips RS. 2000. Experimental erythrocytic malaria infection induces elevated levels of serum amyloid P production in mice. *Immunol Lett* 72:147–152.
- Bayramli G, Ulutas B. 2008. Acute phase protein response in dogs with experimentally induced gastric mucosal injury. *Vet Clin Pathol* 37:312–316.
- Caldin M, Tasca S, Carli E, Bianchini S, Furlanello T, Martinez-Subiela S, Ceron JJ. 2009. Serum acute phase protein concentrations in dogs with hyperadrenocorticism with and without concurrent inflammatory conditions. *Vet Clin Pathol* 38:63–68.
- Castell JV, Andus T, Kunz D, Heinrich PC. 1989. Interleukin 6. The major regulator of acute-phase protein synthesis in man and rat. *Ann N Y Acad Sci* 557:87–99.
- Ceron JJ, Eckersall PD, Martynetz-Subiela S. 2005. Acute phase proteins in dogs and cats: current knowledge and future perspectives. *Vet Clin Pathol* 34:85–99.
- Ceron JJ, Martinez-Subiela S, Ohno K, Caldin M. 2008. A 7-point plan for acute phase protein interpretation in companion animals. *Vet J* 177:6–7.
- Chamanza R, Toussaint MJ, van Ederen AM, van Veen L, Hulskamp-Koch C, Fabri TH. 1999. Serum amyloid A and transferrin in chicken. A preliminary investigation of using acute-phase variables to assess diseases in chickens. *Vet Q* 21:158–162.
- Chamanza R, van Veen L, Tivapasi MT, Toussaint MJ. 1999. Acute phase proteins in the domestic fowl. *Worlds Poult Sci J* 55:61–71.
- Chiarella P, Vulcano M, Bruzzo J, Vermeulen M, Vanzulli S, Maglioco A, Camerano G, Palacios V, Fernandez G, Brando RF, Isturiz MA, Dran GI, Bustuabad OD, Ruggiero RA. 2008. Anti-inflammatory pretreatment enables an efficient dendritic cell-based immunotherapy against established tumors. *Cancer Immunol Immunother* 57:701–718.
- Christou NV, Tellado-Rodriguez J, Chartrand L, Giannas B, Kapadia B, Meakins J, Rode H, Gordon J. 1989. Estimating mortality risk in preoperative patients using immunologic, nutritional, and acute-phase response variables. *Ann Surg* 210:69–77.
- Cohen J, Bayston K. 1990. Lymphokines and the acute-phase response in clinical bone marrow transplantation. *Eur Cytokine Netw* 1:251–255.
- Coles EH. 1974. Liver function, p 203–214. In: Coles EH, editor. *Veterinary clinical pathology*. Philadelphia (PA): WB Saunders.
- Correa SS, Mauldin GN, Mauldin GE, Mooney SC. 2001. Serum alpha 1-acid glycoprotein concentration in cats with lymphoma. *J Am Anim Hosp Assoc* 37:153–158.
- Cray C, Tatum L. 1998. Application of protein electrophoresis in avian diagnostic testing. *J Avian Med Surg* 12:4–10.
- Cray C, Zielzienski-Roberts K, Bonda M, Stevenson R, Ness R, Clubb S, Marsh A. 2005. Antemortem diagnosis of sarcocytosis in psittacine birds: 16 cases. *J Avian Med Surg* 19:208–215.
- Dabrowski R, Wawron W, Kostro K. 2007. Changes in CRP, SAA, and haptoglobin produced in response to ovariectomy in healthy bitches and those with pyometra. *Theriogenology* 67:321–327.
- de Villiers WJ, Varilek GW, de Beer FC, Guo JT, Kindy MS. 2000. Increased serum amyloid A levels reflect colitis severity and precede amyloid formation in IL2-knockout mice. *Cytokine* 12:1337–1347.
- Deak T, Meriwether JL, Fleshner M, Spencer RL, Abouhamze A, Moldawer LL, Grahn RE, Watkins LR, Maier SF. 1997. Evidence that brief stress may induce the acute phase response in rats. *Am J Physiol* 273:R1998–R2004.
- Deans C, Wigmore SJ. 2005. Systemic inflammation, cachexia, and prognosis in patients with cancer. *Curr Opin Clin Nutr Metab Care* 8:265–269.
- Dennis MB Jr. 2000. Humane endpoints for genetically engineered animal models. *ILAR J* 41:94–98.
- Dennis MB Jr. 2002. Welfare issues of genetically modified animals. *ILAR J* 43:100–109.
- Dietsch GN, Dipalma CR, Eyre RJ, Pham TQ, Poole KM, Pefaur NB, Welch WD, Trueblood E, Kerns WD, Kanaly ST. 2006. Characterization of the inflammatory response to a highly selective PDE4 inhibitor in the rat and the identification of biomarkers that correlate with toxicity. *Toxicol Pathol* 34:39–51.
- Duthie S, Eckersall PD, Addie DD, Lawrence CE, Jarrett O. 1997. Value of  $\alpha$ 1-acid glycoprotein in the diagnosis of feline infectious peritonitis. *Vet Rec* 141:299–303.
- Eckersall PD. 1995. Acute phase proteins as markers of inflammatory lesions. *Comp Haematol Int* 5:93–97.
- Eckersall PD. 2000. Recent advances and future prospects for the use of acute phase proteins as markers of disease in animals. *Rev Med Vet (Toulouse)* 151:577–584.
- Eckersall PD, Conner JG, Harvie J. 1991. An immunoturbidimetric assay for canine C-reactive protein. *Vet Res Commun* 15:17–24.
- Eckersall PD, Duthie S, Toussaint MJ, Gruys E, Heegaard P, Alava M, Lipperheide C, Madec F. 1999. Standardization of diagnostic assays for animal acute phase proteins. *Adv Vet Med* 41:643–655.
- Eckersall PD, Gow JW, McComb C, Bradley B, Rodgers J, Murray M, Kennedy PG. 2001. Cytokines and the acute phase response in posttreatment-reactive encephalopathy of *Trypanosoma brucei brucei* infected mice. *Parasitol Int* 50:15–26.
- Eckersall PD, Lawson FP, Bence L, Waterston MM, Lang TL, Donachie W, Fontaine MC. 2007. Acute phase protein response in

- an experimental model of ovine caseous lymphadenitis. *BMC Vet Res* 3:35.
35. **Eckersall PD, Saini PK, McComb C.** 1996. The acute phase response of acid soluble glycoprotein, alpha(1)-acid glycoprotein, ceruloplasmin, haptoglobin, and C-reactive protein in the pig. *Vet Immunol Immunopathol* 51:377-385.
  36. **Endre ZH, Westhuyzen J.** 2008. Early detection of acute kidney injury: emerging new biomarkers. *Nephrology (Carlton)* 13:91-98.
  37. **Forrester S, Hung KE, Kuick R, Kucherlapati R, Haab BB.** 2007. Low-volume, high-throughput sandwich immunoassays for profiling plasma proteins in mice: identification of early-stage systemic inflammation in a mouse model of intestinal cancer. *Mol Oncol* 1:216-225.
  38. **Fransson BA, Bergstrom A, Wardrop KJ, Hagman R.** 2007. Assessment of 3 automated assays for C-reactive protein determination in dogs. *Am J Vet Res* 68:1281-1286.
  39. **French T.** 1989. Acute phase proteins, p 201-235. In: Loeb WF, Quimby FW, editors. *The clinical chemistry of laboratory animals*. Oxford (UK): Pergamon Press.
  40. **Ganheim C, Alenius S, Persson Waller K.** 2007. Acute phase proteins as indicators of calf herd health. *Vet J* 173:645-651.
  41. **Gentry PA, Lumsden JH.** 1978. Determination of serum albumin in domestic animals using the immediate bromocresol green reaction. *Vet Clin Pathol* 7:12-15.
  42. **Giclas PC, Manthei U, Strunk RC.** 1985. The acute phase response of C3, C5, ceruloplasmin, and C-reactive protein induced by turpentine pleurisy in the rabbit. *Am J Pathol* 120:146-156.
  43. **Giffen PS, Turton J, Andrews CM, Barrett P, Clarke CJ, Fung KW, Munday MR, Roman IF, Smyth R, Walshe K, York MJ.** 2003. Markers of experimental acute inflammation in the Wistar Han rat with particular reference to haptoglobin and C-reactive protein. *Arch Toxicol* 77:392-402.
  44. **Giordano A, Spagnolo V, Colombo A, Paltrinieri S.** 2004. Changes in some acute phase protein and immunoglobulin concentrations in cats affected by feline infectious peritonitis or exposed to feline coronavirus infection. *Vet J* 167:38-44.
  45. **Gomez CR, Goral J, Ramirez L, Kopf M, Kovacs EJ.** 2006. Aberrant acute-phase response in aged interleukin-6 knockout mice. *Shock* 25:581-585.
  46. **Grau-Roma L, Heegaard PM, Hjulsager CK, Sibila M, Kristensen CS, Allepuz A, Pineiro M, Larsen LE, Segales J, Fraile L.** 2009. Pig: major acute phase protein and haptoglobin serum concentrations correlate with PCV2 viremia and the clinical course of postweaning multisystemic wasting syndrome. *Vet Microbiol* 138:53-61.
  47. **Gruys E, Toussaint MJ, Niewold TA, Koopmans SJ, van Dijk E, Meloen RH.** 2006. Monitoring health by values of acute phase proteins. *Acta Histochem* 108:229-232.
  48. **Hayashi S, Jinbo T, Iguchi K, Shimizu M, Shimada T, Nomura M, Ishida Y, Yamamoto S.** 2001. A comparison of the concentrations of C-reactive protein and  $\alpha$ 1-acid glycoprotein in the serum of young and adult dogs with acute inflammation. *Vet Res Commun* 25:117-126.
  49. **Hickey MC, Drennan M, Earley B.** 2003. The effect of abrupt weaning of suckler calves on the plasma concentrations of cortisol, catecholamines, leukocytes, acute-phase proteins, and in vitro interferon  $\gamma$  production. *J Anim Sci* 81:2847-2855.
  50. **Hilliquin P.** 1995. Biological markers in inflammatory rheumatic diseases. *Cell Mol Biol (Noisy-Le-Grand)* 41:993-1006.
  51. **Hobo S, Niwa H, Anzai T.** 2007. Evaluation of serum amyloid A and surfactant protein D in sera for identification of the clinical condition of horses with bacterial pneumonia. *J Vet Med Sci* 69:827-830.
  52. **Hol PR, Gruys E.** 1984. Amyloid A proteins in different species. *Appl Pathol* 2:316-327.
  53. **Horadagoda NU, Knox KM, Gibbs HA, Reid SW, Horadagoda A, Edwards SE, Eckersall PD.** 1999. Acute phase proteins in cattle: discrimination between acute and chronic inflammation. *Vet Rec* 144:437-441.
  54. **Hukkanen RR, Liggitt HD, Anderson DM, Kelley ST.** 2006. Detection of systemic amyloidosis in the pigtailed macaque (*Macaca nemestrina*). *Comp Med* 56:119-127.
  55. **Hulten C, Demmers S.** 2002. Serum amyloid A (SAA) as an aid in the management of infectious disease in the foal: comparison with total leucocyte count, neutrophil count, and fibrinogen. *Equine Vet J* 34:693-698.
  56. **Hulten C, Gronlund U, Hirvonen J, Tulamo RM, Suominen MM, Marhaug G, Forsberg M.** 2002. Dynamics in serum of the inflammatory markers serum amyloid A (SAA), haptoglobin, fibrinogen and  $\alpha$ 2-globulins during induced noninfectious arthritis in the horse. *Equine Vet J* 34:699-704.
  57. **Ichiba T, Teshima T, Kuick R, Misek DE, Liu C, Takada Y, Maeda Y, Reddy P, Williams DL, Hanash SM, Ferrara JL.** 2003. Early changes in gene expression profiles of hepatic GVHD uncovered by oligonucleotide microarrays. *Blood* 102:763-771.
  58. **Inoue M, Satoh W, Murakami H.** 1997. Plasma alpha 1-acid glycoprotein in chickens infected with infectious bursal disease virus. *Avian Dis* 41:164-170.
  59. **Jacobsen S, Anderson PH.** 2007. The acute phase protein serum amyloid A (SAA) as a marker of inflammation in horses. *Eq Vet Educ* 19:38-46.
  60. **Jacobsen S, Kjelgaard-Hansen M.** 2008. Evaluation of a commercially available apparatus for measuring the acute phase protein serum amyloid A in horses. *Vet Rec* 163:327-330.
  61. **Jacobsen S, Kjelgaard-Hansen M, Hagbard Petersen H, Jensen AL.** 2006. Evaluation of a commercially available human serum amyloid A (SAA) turbidometric immunoassay for determination of equine SAA concentrations. *Vet J* 172:315-319.
  62. **Jacobsen S, Niewold TA, Halling-Thomsen M, Nanni S, Olsen E, Lindegaard C, Andersen PH.** 2006. Serum amyloid A isoforms in serum and synovial fluid in horses with lipopolysaccharide-induced arthritis. *Vet Immunol Immunopathol* 110:325-330.
  63. **Jacobsen S, Thomsen MH, Nanni S.** 2006. Concentrations of serum amyloid A in serum and synovial fluid from healthy horses and horses with joint disease. *Am J Vet Res* 67:1738-1742.
  64. **Janeway CA, Travers P, Walport M, Shlomchik MJ.** 2001. *Immunobiology*, 5th ed, p732. London (UK): Taylor & Francis.
  65. **Jergens AE, Schreiner CA, Frank DE, Niyo Y, Ahrens FE, Eckersall PD, Benson TJ, Evans R.** 2003. A scoring index for disease activity in canine inflammatory bowel disease. *J Vet Intern Med* 17:291-297.
  66. **Jinbo T, Ami Y, Suzaki Y, Kobune F, Ro S, Naiki M, Iguchi K, Yamamoto S.** 1999. Concentrations of C-reactive protein in normal monkeys (*Macaca irus*) and in monkeys inoculated with *Bordetella bronchiseptica* R5 and measles virus. *Vet Res Commun* 23:265-274.
  67. **Jinbo T, Hayashi S, Iguchi K, Shimizu M, Matsumoto T, Naiki M, Yamamoto S.** 1998. Development of monkey C-reactive protein (CRP) assay methods. *Vet Immunol Immunopathol* 61:195-202.
  68. **Jinbo T, Motoki M, Yamamoto S.** 2001. Variation of serum  $\alpha$ 2-macroglobulin concentration in healthy rats and rats inoculated with *Staphylococcus aureus* or subjected to surgery. *Comp Med* 51:332-335.
  69. **Jinbo T, Sakamoto T, Yamamoto S.** 2002. Serum  $\alpha$ 2-macroglobulin and cytokine measurements in an acute inflammation model in rats. *Lab Anim* 36:153-157.
  70. **Johnson HL, Chiou CC, Cho CT.** 1999. Applications of acute phase reactants in infectious diseases. *J Microbiol Immunol Infect* 32:73-82.
  71. **Kajikawa T, Furuta A, Onishi T, Tajima T, Sugii S.** 1999. Changes in concentrations of serum amyloid A protein,  $\alpha$ 1-acid glycoprotein, haptoglobin, and C-reactive protein in feline sera due to induced inflammation and surgery. *Vet Immunol Immunopathol* 68:91-98.
  72. **Kaneko JJ.** 1997. Serum proteins and the dysproteinemias, p 117-138. In: Kaneko JJ, Harvey JW, Bruss ML, editors. *Clinical biochemistry of domestic animals*. San Diego (CA): Academic Press.
  73. **Kao PC, Shiesh SC, Wu TJ.** 2006. Serum C-reactive protein as a marker for wellness assessment. *Ann Clin Lab Sci* 36:163-169.



74. **Ko KW, Corry DB, Brayton CF, Paul A, Chan L.** 2009. Extravascular inflammation does not increase atherosclerosis in apoE-deficient mice. *Biochem Biophys Res Commun* **384**:93–99.
75. **Lockwood JF, Rutherford MS, Myers MJ, Schook LB.** 1994. Induction of hepatic acute-phase protein transcripts: differential effects of acute and subchronic dimethylnitrosamine exposure in vivo. *Toxicol Appl Pharmacol* **125**:288–295.
76. **Lomborg SR, Nielsen LR, Heegaard PM, Jacobsen S.** 2008. Acute phase proteins in cattle after exposure to complex stress. *Vet Res Commun* **32**:575–582.
77. **MacGuire JG, Christe KL, Yee JL, Kalman-Bowlus AL, Lerche NW.** 2009. Serologic evaluation of clinical and subclinical secondary hepatic amyloidosis in rhesus macaques (*Macaca mulatta*). *Comp Med* **59**:168–173.
78. **Mansfield CS, James FE, Robertson ID.** 2008. Development of a clinical severity index for dogs with acute pancreatitis. *J Am Vet Med Assoc* **233**:936–944.
79. **Martinez-Subiela S, Tecles F, Eckersall PD, Ceron JJ.** 2002. Serum concentrations of acute phase proteins in dogs with leishmaniasis. *Vet Rec* **150**:241–244.
80. **Mischke R, Waterston M, Eckersall PD.** 2007. Changes in C-reactive protein and haptoglobin in dogs with lymphatic neoplasia. *Vet J* **174**:188–192.
81. **Murata H.** 2007. Stress and acute phase protein response: an inconspicuous but essential linkage. *Vet J* **173**:473–474.
82. **Murata H, Shimada N, Yoshioka M.** 2004. Current research on acute phase proteins in veterinary diagnosis: an overview. *Vet J* **168**:28–40.
83. **Myers LA, Boyce JT, Robison RL.** 1995. The tolerability and pharmacology of interleukin 6 administered in combination with GM-CSF or G-CSF in the rhesus monkey. *Toxicology* **101**:157–166.
84. **Nakamura K, Mitarai Y, Yoshioka M, Koizumi N, Shibahara T, Nakajima Y.** 1998. Serum levels of interleukin 6,  $\alpha$ 1-acid glycoprotein, and corticosterone in 2-week-old chickens inoculated with *Escherichia coli* lipopolysaccharide. *Poult Sci* **77**:908–911.
85. **Nakamura M, Takahashi M, Ohno K, Koshino A, Nakashima K, Setoguchi A, Fujino Y, Tsujimoto H.** 2008. C-reactive protein concentration in dogs with various diseases. *J Vet Med Sci* **70**:127–131.
86. **Nemzek JA, Xiao HY, Minard AE, Bolgos GL, Remick DG.** 2004. Humane endpoints in shock research. *Shock* **21**:17–25.
87. **Nielsen L, Toft N, Eckersall PD, Mellor DJ, Morris JS.** 2007. Serum C-reactive protein concentration as an indicator of remission status in dogs with multicentric lymphoma. *J Vet Intern Med* **21**:1231–1236.
88. **Ohno K, Yokoyama Y, Nakashima K, Setoguchi A, Fujino Y, Tsujimoto H.** 2006. C-reactive protein concentration in canine idiopathic polyarthritis. *J Vet Med Sci* **68**:1275–1279.
89. **Olfert ED, Godson DL.** 2000. Humane endpoints for infectious disease animal models. *ILAR J* **41**:99–104.
90. **Paltrinieri S.** 2008. The feline acute phase reaction. *Vet J* **177**:26–35.
91. **Parra MD, Fuentes P, Tecles F, Martinez-Subiela S, Martinez JS, Munoz A, Ceron JJ.** 2006. Porcine acute phase protein concentrations in different diseases in field conditions. *J Vet Med B Infect Dis Vet Public Health* **53**:488–493.
92. **Pepys MB, Hirschfield GM.** 2003. C-reactive protein: a critical update. *J Clin Invest* **111**:1805–1812.
93. **Petersen HH, Nielsen JP, Heegaard PM.** 2004. Application of acute phase protein measurements in veterinary clinical chemistry. *Vet Res* **35**:163–187.
94. **Pfeffer A, Rogers KM.** 1989. Acute phase response of sheep: changes in the concentrations of ceruloplasmin, fibrinogen, haptoglobin, and the major blood cell types associated with pulmonary damage. *Res Vet Sci* **46**:118–124.
95. **Pfeffer A, Rogers KM, O'Keeffe L, Osborn PJ.** 1993. Acute phase protein response, food intake, live-weight change, and lesions following intrathoracic injection of yeast in sheep. *Res Vet Sci* **55**:360–366.
96. **Pineiro M, Pineiro C, Carpintero R, Morales J, Campbell FM, Eckersall PD, Toussaint MJ, Lampreave F.** 2007. Characterisation of the pig acute phase protein response to road transport. *Vet J* **173**:669–674.
97. **Pollock PJ, Prendergast M, Schumacher J, Bellenger CR.** 2005. Effects of surgery on the acute phase response in clinically normal and diseased horses. *Vet Rec* **156**:538–542.
98. **Pruett BS, Pruett SB.** 2006. An explanation for the paradoxical induction and suppression of an acute phase response by ethanol. *Alcohol* **39**:105–110.
99. **Quinton LJ, Jones MR, Robson BE, Mizgerd JP.** 2009. Mechanisms of the hepatic acute-phase response during bacterial pneumonia. *Infect Immun* **77**:2417–2426.
100. **Raida MK, Buchmann K.** 2009. Innate immune response in rainbow trout (*Oncorhynchus mykiss*) against primary and secondary infections with *Yersinia ruckeri* O1. *Dev Comp Immunol* **33**:35–45.
101. **Ray A, Ray BK.** 1999. Persistent expression of serum amyloid A during experimentally induced chronic inflammatory condition in rabbit involves differential activation of SAF, NF $\kappa$ B, and CEBP transcription factors. *J Immunol* **163**:2143–2150.
102. **Ridker PM.** 2007. Inflammatory biomarkers and risks of myocardial infarction, stroke, diabetes, and total mortality: implications for longevity. *Nutr Rev* **65**:S253–S259.
103. **Rush JE, Lee ND, Freeman LM, Brewer B.** 2006. C-reactive protein concentration in dogs with chronic valvular disease. *J Vet Intern Med* **20**:635–639.
104. **Saco Y, Fina M, Gimenez M, Pato R, Piedrafita J, Bassols A.** 2008. Evaluation of serum cortisol, metabolic parameters, acute phase proteins and faecal corticosterone as indicators of stress in cows. *Vet J* **177**:439–441.
105. **Saini PK, Riaz M, Webert DW, Eckersall PD, Young CR, Stanker LH, Chakrabarti E, Judkins JC.** 1998. Development of a simple enzyme immunoassay for blood haptoglobin concentration in cattle and its application in improving food safety. *Am J Vet Res* **59**:1101–1107.
106. **Salamano G, Mellia E, Candiani D, Ingravalle F, Bruno R, Ru G, Doglione L.** 2008. Changes in haptoglobin, C-reactive protein and pig-major acute phase protein (porcine) during a housing period following long distance transport in swine. *Vet J* **177**:110–115.
107. **Sandor G.** 1966. Serum proteins in health and disease. Baltimore (MD): Williams and Wilkins.
108. **Schreiber G, Tsykin A, Aldred AR, Thomas T, Fung WP, Dickson PW, Cole T, Birch H, De Jong FA, Milland J.** 1989. The acute phase response in the rodent. *Ann N Y Acad Sci* **557**:61–85.
109. **Selting KA, Ogilvie GK, Lana SE, Fettman MJ, Mitchener KL, Hansen RA, Richardson KL, Walton JA, Scherk MA.** 2000. Serum  $\alpha$ 1-acid glycoprotein concentrations in healthy and tumor-bearing cats. *J Vet Intern Med* **14**:503–506.
110. **Shimada T, Ishida Y, Shimizu M, Nomura M, Kawato K, Iguchi K, Jinbo T.** 2002. Monitoring C-reactive protein in beagle dogs experimentally inoculated with *Ehrlichia canis*. *Vet Res Commun* **26**:171–177.
111. **Skinner JG.** 2001. International standardization of acute phase proteins. *Vet Clin Pathol* **30**:2–7.
112. **Skinner JG, Roberts L.** 1994. Haptoglobin as an indicator of infection in sheep. *Vet Rec* **134**:33–36.
113. **Skovgaard K, Mortensen S, Boye M, Poulsen KT, Campbell FM, Eckersall PD, Heegaard PM.** 2009. Rapid and widely disseminated acute phase protein response after experimental bacterial infection of pigs. *Vet Res* **40**:23.
114. **Solter PF, Hoffmann WE, Hungerford LL, Siegel JP, St Denis SH, Dorner JL.** 1991. Haptoglobin and ceruloplasmin as determinants of inflammation in dogs. *Am J Vet Res* **52**:1738–1742.
115. **Stokol T, Tarrant JM, Scarlett JM.** 2001. Overestimation of canine albumin concentration with the bromocresol green method in heparinized plasma samples. *Vet Clin Pathol* **30**:170–176.
116. **Suffredini AF, Fantuzzi G, Badolato R, Oppenheim JJ, O'Grady NP.** 1999. New insights into the biology of the acute phase response. *J Clin Immunol* **19**:203–214.

117. **Takahashi K, Miyake N, Ohta T, Akiba Y, Tamura K.** 1998. Changes in plasma  $\alpha$ 1-acid glycoprotein concentration and selected immune response in broiler chickens injected with *Escherichia coli* lipopolysaccharide. *Br Poult Sci* **39**:152–155.
118. **Tamamoto T, Ohno K, Ohmi A, Goto-Koshino Y, Tsujimoto H.** 2008. Verification of measurement of the feline serum amyloid A (SAA) concentration by human SAA turbidimetric immunoassay and its clinical application. *J Vet Med Sci* **70**:1247–1252.
119. **Tamamoto T, Ohno K, Ohmi A, Seki I, Tsujimoto H.** 2009. Time-course monitoring of serum amyloid A in a cat with pancreatitis. *Vet Clin Pathol* **38**:83–86.
120. **Tecles F, Caldin M, Zanella A, Membiela F, Tvarijonaviciute A, Subiela SM, Ceron JJ.** 2009. Serum acute phase protein concentrations in female dogs with mammary tumors. *J Vet Diagn Invest* **21**:214–219.
121. **Tecles F, Spiranelli E, Bonfanti U, Ceron JJ, Paltrinieri S.** 2005. Preliminary studies of serum acute-phase protein concentrations in hematologic and neoplastic diseases of the dog. *J Vet Intern Med* **19**:865–870.
122. **Tecles F, Subiela SM, Petrucci G, Panizo CG, Ceron JJ.** 2007. Validation of a commercially available human immunoturbidimetric assay for haptoglobin determination in canine serum samples. *Vet Res Commun* **31**:23–36.
123. **Tillett WS, Francis T.** 1930. Serological reactions in pneumonia with a nonprotein somatic fraction of *Pneumococcus*. *J Exp Med* **52**:561–571.
124. **Tohjo H, Yadatsu M, Uchida E, Niiyama M, Syuto B, Moritsu Y, Ichikawa S, Takeuchi M.** 1996. Polyacrylamide gel electrophoretic serum protein patterns of acute inflammation induced by intramuscular injection of turpentine in young broiler chickens. *J Vet Med Sci* **58**:267–268.
125. **Toth LA.** 2000. Defining the moribund condition as an experimental endpoint for animal research. *ILAR J* **41**:72–79.
126. **Toussaint MJ, van Ederen AM, Gruys E.** 1995. Implication of clinical pathology in assessment of animal health and in animal production and meat inspection. *Comp Haematol Int* **5**:149–157.
127. **Ulutas B, Bayramli G, Ulutas PA, Karagenc T.** 2005. Serum concentration of some acute phase proteins in naturally occurring canine babesiosis: a preliminary study. *Vet Clin Pathol* **34**:144–147.
128. **Ulutas PA, Ozpinar A.** 2006. Effect of *Mannheimia (Pasteurella) haemolytica* infection on acute-phase proteins and some mineral levels in colostrum–breast-milk-fed or colostrum–breast-milk-deprived sheep. *Vet Res Commun* **30**:485–495.
129. **Van Gool J, Van Vugt H, Helle M, Aarden LA.** 1990. The relation among stress, adrenalin, interleukin 6, and acute phase proteins in the rat. *Clin Immunol Immunopathol* **57**:200–210.
130. **Vandenplas ML, Moore JN, Barton MH, Roussel AJ, Cohen ND.** 2005. Concentrations of serum amyloid A and lipopolysaccharide-binding protein in horses with colic. *Am J Vet Res* **66**:1509–1516.
131. **Vernooy JH, Reynaert N, Wolfs TG, Cloots RH, Haegens A, de Vries B, Dentener MA, Buurman WA, Wouters EM.** 2005. Rapid pulmonary expression of acute-phase reactants after local lipopolysaccharide exposure in mice is followed by an interleukin-6-mediated systemic acute-phase response. *Exp Lung Res* **31**:855–871.
132. **Xie H, Newberry L, Clark FD, Huff WE, Huff GR, Balog JM, Rath NC.** 2002. Changes in serum ovotransferrin levels in chickens with experimentally induced inflammation and diseases. *Avian Dis* **46**:122–131.
133. **Yamamoto S, Shida T, Miyaji S, Santsuka H, Fujise H, Mukawa K, Furukawa E, Nagae T, Naiki M.** 1993. Changes in serum C-reactive protein levels in dogs with various disorders and surgical traumas. *Vet Res Commun* **17**:85–93.
134. **Yamamoto S, Shida T, Okimura T, Otabe K, Honda M, Ashida Y, Furukawa E, Sarikaputi M, Naiki M.** 1994. Determination of C-reactive protein in serum and plasma from healthy dogs and dogs with pneumonia by ELISA and slide reversed passive latex agglutination test. *Vet Q* **16**:74–77.
135. **Yamamoto S, Tagata K, Nagahata H, Ishikawa Y, Morimatsu M, Naiki M.** 1992. Isolation of canine C-reactive protein and characterization of its properties. *Vet Immunol Immunopathol* **30**:329–339.
136. **Yamashita K, Fujinaga T, Miyamoto T, Hagio M, Izumisawa Y, Kotani T.** 1994. Canine acute phase response: relationship between serum cytokine activity and acute phase protein in dogs. *J Vet Med Sci* **56**:487–492.
137. **Zaias J, Mineau M, Cray C, Yoon D, Altman NH.** 2009. Baseline reference values for serum protein electrophoresis in common laboratory rodent strains. *J Am Assoc Lab Anim Sci* **48**:387–390.