

Original Research

Subregional Expression of Hippocampal Glutamatergic and GABAergic Genes in F344 Rats with Social Isolation after Weaning

Hisaya Iwata and Yutaka Yamamuro*

Many studies have shown that postweaning social isolation (pwSI) alters various behavioral phenotypes, including hippocampus-dependent tasks. Here, we report the comprehensive analysis of the expression of glutamatergic and GABAergic neurotransmission-related genes in the distinct hippocampal subregions of pwSI rats. Male F344 rats (age, 4 wk) experienced either pwSI or group housing (controls). At 7 wk of age, the hippocampus of each rat was removed and laser-microdissected into the CA1 and CA3 layers of pyramidal cells and the granule cell layer of the dentate gyrus. Subsequently, the expression of glutamatergic- and GABAergic-related genes was analyzed by quantitative RT-PCR. In the CA1 and CA3 pyramidal cell layers, 18 of 24 glutamate receptor subunit genes were at least 1.5-fold increased in expression after pwSI. In particular, the expression of several N-methyl-D-aspartate and kainate receptors (for example, *Grin2a* in CA1, *Grik4* in CA3) was significantly increased after pwSI. In contrast, pwSI tended to decrease the expression of GABA_A receptor subunit genes, and *Gabra1*, *Gabra2*, *Gabra4*, *Gabra5*, *Gabrb2*, *Gabrg1*, and *Gabrg2* were all significantly decreased in expression compared with the levels in the group-housed rats. These results indicate a subregion-specific increase of glutamate receptors and reduction of GABA_A receptors, suggesting that the hippocampal circuits of pwSI rats may be in more excitable states than those of group-housed rats.

Abbreviations: AMPA, α -amino-2,3-dihydro-5-methyl-3-oxo-4-isoxazolepropanoic acid; LMD, laser microdissection; NMDA; N-methyl-D-aspartic acid; pwSI, postweaning social isolation; DG; dentate gyrus

Behavioral phenotypes differ among strains of laboratory animals, including mice and rats. The acquisition of strain-specific behavioral traits is governed not only by the genetic background but also by the postnatal rearing environment. The level of maternal care is a representative environmental factor. For example, inbred Fischer 344 pups raised by Wistar dams showed Wistar-like behavioral traits in adulthood in socially interactive situations.³⁵ Furthermore, the effect of environmental factors is not limited to neonatal periods. In animals that form a society among conspecifics, such as mice and rats, postweaning social environments have a marked effect on behavioral traits. Many studies have reported that postweaning social isolation (pwSI) alters various strain-specific behavioral phenotypes, including aggressiveness,^{15,32,34} novelty preference,²² locomotor activity,²⁹ anxiety-like behavior,²⁰ and learning and memory.^{13,15,20,31} In other words, pwSI means the deprivation of several social interactions among conspecifics. The expression of social contact begins fundamentally in the postweaning periods, its frequency increases to a peak at 4 to 5 wk of age, and it declines thereafter until sexual maturity.^{2,17,30} Therefore, rats were isolated from their conspecifics to deprive them of social contact during the current study.

In the brain, the hippocampus is necessary for the acquisition of episodic and spatial memory,²⁸ but the influences of pwSI on hippocampal functions remain largely unclear. The hippocampus has a lamellar organization of neurons, and the intrinsic neuronal circuit of the hippocampus, termed the trisynaptic circuit, consists of 3 topographically and morphologically distinct neuronal layers: the pyramidal cell layer in subfields CA1 and CA3 and the granule cell layer in the dentate gyrus (DG). Sensory information is carried first to the DG by a perforant pathway that originates in the entorhinal cortex. DG granule cells project to the apical dendrites of the CA3 pyramidal cells through mossy fibers. In turn, CA3 pyramidal cells project to the CA1 layer through Schaffer collaterals.^{1,18} Glutamate is a key excitatory neurotransmitter in the hippocampus and plays a central role in the activation of the trisynaptic circuit, whereas the inhibitory neurotransmitter GABA modulates the activated circuit. This excitatory–inhibitory balance is critical for the appropriate functioning of the hippocampal circuit. Here, we comprehensively investigated the effect of pwSI on the expression of glutamatergic and GABAergic neurotransmission-related genes in the 3 hippocampal subregions of inbred F344 rats.

Materials and Methods

All experiments were performed in accordance with the Guidelines for Animal Experiments (College of Bioresource Sciences,

Received: 14 May 2015. Revision requested: 28 Jun 2015. Accepted: 05 Aug 2015.
Laboratory of Animal Genetics and Physiology, Department of Animal Science, College of Bioresource Sciences, Nihon University, Fujisawa, Kanagawa, Japan
*Corresponding author. Email: yamamuro.yutaka@nihon-u.ac.jp

Nihon University) and Nihon University's Animal Care and Use Committees. Male F344/DuCrIj rats (age, 3 wk; SPF, viral antigen-free) were purchased from Charles River Laboratories Japan (Yokohama, Japan). Rats were housed in opaque polypropylene cages (345 × 403 × 177 mm) and allowed free access to food (Labo MR Stock, Nosan, Kanagawa, Japan) and water. The rats were housed in an animal room maintained at 22 ± 1 °C with a humidity of 55 ± 10% and a 14:10-h light:dark cycle (lights on, 0600). When rats had reached 4 wk of age, they were allocated to 1 of 2 rearing conditions: isolated (pwSI; 1 rat per cage, $n = 3$) or group-housed (3 rats / cage, $n = 3$, with 1 rat randomly obtained from each cage) rearing. Rats were euthanized at 7 wk of age by CO₂, and whole brains were removed. The trimmed brains were embedded in OCT compound and stored at -80 °C.

To separate the 3 hippocampal subregions, we used a laser microdissection system (LMD; model LMD7000-4, Leica Microsystems, Tokyo, Japan). First, coronal sections including the hippocampus (30 μm) were cut by using a cryostat (HM550; Thermo Fischer Scientific, Yokohama, Japan) at -10 °C and mounted on membrane slides (Membrane Slides Nuclease and Human Nucleic Acid Free PEN-Membrane, Leica Microsystems). According to the manufacturer's (Leica Microsystems) recommendations, the sections were fixed in a mixture of acetate and ethanol (1:19, v/v), and briefly stained with toluidine blue. The slides were air-dried and set within the LMD system. The CA1 and CA3 pyramidal cell layers and DG granule cell layers were cut according to the Allen Brain Atlas (<http://mouse.brain-map.org/static/atlas>) and collected separately into the caps of 0.2-mL microtubes (BT-02LC; Ina-Optica, Osaka, Japan) supplied with the LMD system.

The quantitative RT-PCR protocol was modified from previous methods.³⁶ The collected specimens were completely dissolved in 9.5 mL of the lysis buffer provided with CellAmp Direct RNA Prep Kit (Takara Bio, Ohtsu, Japan) supplemented with proteinase K (0.3 U, Takara Bio). DNA in the lysate was digested by adding 1 mL of DNase provided with the kit and incubating at 37 °C for 5 min. Then, distilled water (UltraPure DNase/RNase-Free Distilled Water; Life Technologies, Carlsbad, CA) was added to the lysate, which was incubated at 75 °C for 5 min to inactivate DNase. The cDNA was synthesized by using the iScript cDNA Synthesis Kit (BioRad, Hercules, CA) according to the manufacturer's instructions. Real-time PCR was performed by using Mini Opticon (BioRad) with SsoFast EvaGreen qPCR SuperMix (BioRad). Gene-specific primers were designed by using Primer Blast (<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>), and all primer sequences are listed in Table 1.

Statistical significance was determined by using 2-way ANOVA, and differences between control and pwSI groups when comparing the same region were tested in subsequent planned comparisons, contrasting the mean comparisons of selected levels of a factor, by using the software package Super ANOVA (Abacus Concepts, Berkeley, CA). The criterion for significance was a P value less than 0.05 in all cases.

Results

The glutamatergic neurotransmission-related genes measured in the present study were α -amino-2,3-dihydro-5-methyl-3-oxo-4-isoxazolepropanoic acid (AMPA) receptor subunits (*Gria1–Gria4*), NMDA receptor subunits (*Grin1–Grin3b*), kainate receptor subunits (*Grik1–Grik5*), metabotropic receptor subunits (*Grm1–Grm8*),

vesicular glutamate transporters (*Slc17a6–Slc17a8*), membrane glutamate transporters (*Slc1a1, Slc1a6*), and glutamate synthases (*Gls, Gls2*). These results are shown in Table 2.

In CA1 and CA3 pyramidal cell layers, 18 of the 24 glutamate receptor subunit genes evaluated were at least 1.5-fold increased ($P < 0.05$) in expression in pwSI rats compared with group-housed rats, and only *Gria3* was decreased significantly ($P < 0.05$) by pwSI. In particular, *Grin2a* ($P = 0.0003$) in CA1 and *Grik4* ($P = 0.0065$) in CA3 were significantly ($P < 0.05$) increased by pwSI. In contrast, pwSI had little effect on the expression of the glutamate receptor subunits in the DG granule cell layer, whereas the vesicular transporters, membrane transporters, and synthases tended to increase. The GABAergic neurotransmission-related genes analyzed were: GABA_A receptor α subunits (*Gabra1–Gabra6*), β subunits (*Gabrb1–Gabrb3*), γ subunits (*Gabrg1–Gabrg3*), δ subunit (*Gabrd*), GABA_B receptor subunits (*Gabbr1, Gabbr2*), vesicular GABA transporter (*Slc32a1*), membrane GABA transporter (*Slc6a1*), and GABA synthases (*Gad1, Gad2*), and the results are shown in Table 3. Compared with group housing, pwSI had little effect on the expressions of these genes in the CA1 and CA3 regions. In contrast to its effect in the pyramidal cell layers, pwSI tended to decrease the expression of 8 of the 13 GABA_A receptor subunit genes, especially 4 of the 5 α -subunit genes and 3 of 3 γ -subunit genes. Among them, the expression of *Gabra1, Gabra2, Gabra4, Gabra5, Gabrb3, Gabrg1*, and *Gabrg2* was significantly ($P < 0.05$) lower in the DG of pwSI rats than in their group-housed counterparts.

Discussion

The present study showed that the gene expression of GABA_A receptor subunits, except for the GABA_B receptor, in the DG granule cell layer of pwSI rats largely tended to be lower than that in group-housed rats. The GABA_A receptor is a ligand-operated chloride channel composed of 5 subunits. Sixteen subunit isoforms have been identified in mammalian brains:²⁶ 6 α subunits, 3 β subunits, 3 γ subunits, and single δ , ϵ , θ , and π subunits. The predominant GABA_A receptors consist of 2 α , 2 β , and a single γ or δ subunit.^{25,31} The physiologic properties and intracellular localization of GABA_A receptors are governed by the subunit composition of the pentamer.²⁴ For example, GABA_A receptors containing $\alpha 1$ subunits, which were reduced in expression by pwSI in the present study, show postsynaptic localization and mediate phasic synaptic inhibition.¹² In contrast, the receptors that contain $\alpha 4$ or $\alpha 5$, which also showed decreased expression in pwSI rats, are extrasynaptic and induce tonic conductance.¹⁰ Irrespective of these effects, pwSI might reduce the expression of functional GABA_A receptors in the DG granule cell layer. Moreover, our finding that the $\alpha 6$ subunit was not expressed in either the pyramidal cell layer or the granule cell layer is consistent with a previous report.²⁰ Several reports describe the functional consequences of the various GABA_A receptors. A study using sections from adult rat brain revealed that GABA_A receptors containing the $\alpha 5$ subunit are the predominant extrasynaptic inhibitory receptor.³ Rendering this receptor dysfunctional rescues the learning impairment induced by anesthetics⁵ and enhances the learning ability analyzed by a variety of behavioral tasks, including fear conditioning.^{4,6,9} Therefore, the functional consequence of GABA_A receptors continues to become clear.

The expression of NMDA and kainate receptors, rather than AMPA receptors, tended to increase in CA1 and CA3 pyramidal

Table 1. Sequences of the primers used in quantitative RT-PCR

Primer	Forward (5'→3')	Reverse (5'→3')	Accession no.
<i>Gria1</i>	CTTCATGCAGCAAGGATGTG	TGGCTGTGTACGAGGAGATG	NM_031608
<i>Gria2</i>	AGGTGGCAAAGAATCCACAG	GCCTTGCTCCTCATGACATC	NM_017261
<i>Gria3</i>	GGGAAGCATTCAAGAGGCTA	CTGCTGAACCAATTGGGTTTT	NM_032990
<i>Gria4</i>	TTTGCAGGCAGATTGTCTTG	TGGTCTGTGTTTCGGATGAA	NM_017263
<i>Grin1</i>	TCCAGCTTCAAGAGACGTAGG	CTCTCCCTATGACGGGAACA	NM_017010
<i>Grin2a</i>	CTGGAAGAGGCAGATTGACC	TCTTCTCGTTGTGGCAGATG	NM_012573
<i>Grin2b</i>	GCATGCCTACATGGGAAAGT	GTTGAGCACAGCTGCATCAT	NM_012574
<i>Grin2c</i>	GACTTCTGCTGGCTTTTCAG	CTGCAGCATCTTCAGCACAT	NM_012575
<i>Grin2d</i>	TCTTCGCCACCACTGGTTAT	ATTGTGGCAGATCCCTGAAA	NM_022797
<i>Grin3a</i>	GAAGAAAAGCAGCCACGTTT	GGTTTTGTCTTCCTCGTCA	NM_138546
<i>Grin3b</i>	AGGACCTGGCCTTTGACTTT	TTGATGCTGAAGCTGGTCAC	NM_133308
<i>Grik1</i>	TGGCAAATATGGAGCACAGA	GTCATGAAGGGCTTGGAGAA	NM_017241
<i>Grik2</i>	TGGAGGATGGGAAATATGGA	CGATGACCTTCTCACGAACA	NM_019309
<i>Grik3</i>	CGGAGTCTGGTTTGGGAATA	CGTCCGCGTACTCAAAGATT	NM_181373
<i>Grik4</i>	GCTCCAGCATGACCTTCTTC	CCCTCCTCTGTGCTTTCAC	NM_012572
<i>Grik5</i>	TTCTCCTTCTGGACCCTTT	GAAGACACGGGTGAGGGTTA	NM_031508
<i>Grm1</i>	GGTCCCTTCTGACACTTTGC	CATTCCACTCTCGCCGTAAT	NM_017011
<i>Grm2</i>	GGTAACTACGGGGACAGAA	TCACCAGCCCCTGATGTAA	NM_001105711
<i>Grm3</i>	CCAAGTCTGTGATGCAATG	CATCCCGTCTCCAAAAGTGT	NM_001105712
<i>Grm4</i>	TCAAGAAGGGAAGCCACATC	CACAGCGTCAATCACGAACT	NM_022666
<i>Grm5</i>	GTTTGCACAGGAGAACAGCA	AGGGACATCTGCATATTGTGG	NM_017012
<i>Grm6</i>	GAGGCCTTTGTGCAGATTTC	CCGGATCACTTGTGGAACCT	NM_022920
<i>Grm7</i>	CCAGGGCGTTATGACATCTT	AGGATGGGATCTCTCGACT	NM_031040
<i>Grm8</i>	CGAGCAAGGGAAGAAATCTG	CCACGACAAACCACACAAAC	NM_022202
<i>Slc17a6</i>	TCAACCAGCTTATTCGAGGA	ATTGTCATGACCAGGTGTGG	NM_053427
<i>Slc17a7</i>	CTGGTGGTCGGATACTACA	ATTGGAAATGCCCATCAAGA	NM_053859
<i>Slc17a8</i>	ACGGGATGTGGAGTAAGTGG	TACTCCTGCAAGGGGCATAG	NM_153725
<i>Slc1a1</i>	TTCATAGTCGTCCGAAGAA	CACAGCGGAATGTAAGTGA	NM_013032
<i>Slc1a6</i>	CATGCTGGTGGTGTGTTACC	AAGAGGTACCCATGGCTGTG	NM_032065
<i>Gls</i>	CTTGACAAGATGGGCAACAG	TCCTTCTCTCCGAGGATCAA	NM_012569
<i>Gls2</i>	AACAAGACCGTGGTGAACCT	TGCGGGAATCATAGTCCTTC	NM_138904
<i>Gabra1</i>	TCTGAGCACTCTCTCGGAA	AGTTACACGCTCTCCCAAGC	NM_183326
<i>Gabra2</i>	CGTATGGTTTCCGCTGCTTG	GTAATGCTGTCTCCCAGTCCT	NM_001135779
<i>Gabra3</i>	CCAAGGGGAGTCAAGACGAC	AGTCACTGCATCTCCAAGCC	NM_017069
<i>Gabra4</i>	TGTGCCTGGCGACTTGTTA	CCAAATCCAGGACGCAGTCT	NM_080587
<i>Gabra5</i>	TGTCCGACACGGAATGGAAT	TGTTGAGAGGGAGACGTTGC	NM_017295
<i>Gabra6</i>	AACGCTGACTGTCCGATGAG	TCGGATAGGCATAGCTCCA	NM_021841
<i>Gabrb1</i>	ACAACAGGGGCATATCCACG	CAGTGTGGAGGGCATGTAGG	NM_012956
<i>Gabrb2</i>	TGTCAACAAGATGGACCCACA	ATGCTGGAGGCATCATAGGC	NM_012957
<i>Gabrb3</i>	CTGTACGGGCTCAGGATCAC	CGATCCTTTCCACGCCAGTA	NM_017065
<i>Gabrg1</i>	AAACATGGGTCTTGGCACCT	ATTACTGTGGGTCTCACGCC	NM_080586
<i>Gabrg2</i>	GAAAAACCCTGCCCTACCA	TGCGAATGTGTATCCTCCCG	NM_183327
<i>Gabrg3</i>	GAGGCTCACTGGATACCAC	GGGGTAGCCATAGCTAGAGA	NM_024370
<i>Gabrd</i>	GCGCCAGAGCAATGAATGAC	GCGTAGCCCTCCATTAGTCC	NM_017289
<i>Gabbr1</i>	CTCTCGGGCTGGATGGTTAC	GGCTCTAGGGTCTTCCTCCA	NM_031028
<i>Gabbr2</i>	CAGGGAAGACTCCACAGCAG	AGGCGTACCCATGGAACCTG	NM_031802
<i>Slc32a1</i>	CCGTCGAGGGAGACATTCATT	CACGAACATGCCCTGAATGG	NM_031782
<i>Slc6a1</i>	ACAGCCAGTTCTGTACCGTG	CACACGGCAGCAATGAAGAG	NM_024371
<i>Gad1</i>	GATGGTTTTCGATGGT	CCATGGTTGTTCTCTGA	NM_008077
<i>Gad2</i>	GTGTTTCGATGGGAAGCCTCA	TAATCACTGGCGCCACCTTT	NM_012563
<i>Gapdh</i>	ATGACTCTACCCACGGCAAG	TACTCAGCACCAGCATCACC	NM_017008

Table 2. Effect of pwSI on glutamatergic neuron-related gene expressions in distinct hippocampal regions of F344 rats

	CA1		CA3		DG	
	Group	pwSI	Group	pwSI	Group	pwSI
<i>Gria1</i>	1.7E+00 ± 3.0E-01	1.8E+00 ± 3.0E-01	5.3E-01 ± 4.9E-02	7.5E-01 ± 1.2E-02	1.6E+00 ± 2.7E-01	1.9E+00 ± 3.5E-01
<i>Gria2</i>	6.1E-02 ± 1.5E-02	5.9E-02 ± 8.6E-03	1.7E-02 ± 1.3E-03	1.9E-02 ± 5.7E-04	7.2E-02 ± 5.6E-03	5.5E-02 ± 5.7E-03
<i>Gria3</i>	4.8E-02 ± 9.5E-03	3.1E-02 ± 7.1E-03 ^a	1.2E-02 ± 1.1E-03	1.1E-02 ± 4.9E-04	3.4E-02 ± 5.7E-03	4.3E-02 ± 4.1E-03
<i>Gria4</i>	3.4E-01 ± 7.5E-02	4.7E-01 ± 8.8E-02	9.8E-02 ± 1.1E-02	1.6E-01 ± 2.2E-02	6.4E-01 ± 1.7E-01	8.1E-01 ± 1.6E-01
<i>Grin1</i>	6.6E-02 ± 1.4E-02	8.7E-02 ± 1.9E-02	3.4E-02 ± 5.6E-03	4.7E-02 ± 1.5E-03	6.9E-02 ± 1.0E-02	6.7E-02 ± 4.7E-03
<i>Grin2a</i>	3.0E+00 ± 1.4E-02	5.0E+00 ± 4.7E-01^c	9.6E-01 ± 1.6E-01	1.5E+00 ± 8.4E-02	2.5E+00 ± 3.4E-01	2.4E+00 ± 3.3E-01
<i>Grin2b</i>	3.1E+00 ± 7.8E-01	3.9E+00 ± 7.3E-01	1.1E+00 ± 4.9E-02	1.7E+00 ± 2.5E-01	3.4E+00 ± 4.1E-01	3.7E+00 ± 2.5E-01
<i>Grin2c</i>	9.2E-02 ± 1.6E-02	1.1E-01 ± 1.5E-02	3.6E-02 ± 6.4E-03	4.3E-02 ± 5.1E-03	6.4E-02 ± 8.2E-03	8.4E-02 ± 2.4E-02
<i>Grin2d</i>	5.1E-03 ± 4.7E-04	9.1E-03 ± 1.6E-03	3.3E-03 ± 4.1E-04	3.3E-03 ± 4.0E-04	1.3E-02 ± 2.7E-03	1.2E-02 ± 1.4E-03
<i>Grin3a</i>	9.0E-03 ± 2.7E-03	1.3E-02 ± 1.3E-03	2.5E-03 ± 3.8E-04	2.6E-03 ± 2.3E-04	<i>1.0E-02 ± 1.5E-03</i>	<i>6.3E-03 ± 4.7E-04</i>
<i>Grin3b</i>	nd	nd	nd	nd	nd	nd
<i>Grik1</i>	1.6E-01 ± 7.1E-02	2.8E-01 ± 7.6E-02	6.8E-02 ± 5.6E-03	1.9E-01 ± 2.7E-02	4.0E-01 ± 7.6E-02	5.9E-01 ± 8.6E-02
<i>Grik2</i>	4.9E-01 ± 8.0E-02	7.5E-01 ± 9.4E-02	4.9E-01 ± 7.4E-02	8.5E-01 ± 2.0E-01	2.6E+00 ± 3.8E-01	2.6E+00 ± 3.8E-01
<i>Grik3</i>	5.3E-04 ± 1.2E-04	6.3E-04 ± 1.2E-04	2.9E-04 ± 1.8E-05	3.7E-04 ± 4.2E-05	2.9E-03 ± 3.7E-04	2.8E-03 ± 3.5E-04
<i>Grik4</i>	4.7E-02 ± 1.7E-02	4.6E-02 ± 5.3E-03	1.1E+00 ± 9.4E-02	1.7E+00 ± 2.2E-01^b	7.3E-01 ± 1.6E-01	1.0E+00 ± 2.6E-01
<i>Grik5</i>	1.1E+01 ± 2.2E+00	1.3E+01 ± 2.3E+00	6.5E+00 ± 2.8E-01	1.0E+01 ± 1.0E+00	1.3E+01 ± 2.0E+00	1.7E+01 ± 1.4E+00 ^a
<i>Grm1</i>	1.4E-03 ± 2.7E-04	2.3E-03 ± 3.7E-04	1.3E-02 ± 1.4E-04	2.2E-02 ± 3.1E-03	6.4E-02 ± 7.2E-03	7.9E-02 ± 1.6E-03 ^a
<i>Grm2</i>	nd	nd	nd	nd	9.2E-03 ± 1.1E-03	7.6E-03 ± 3.0E-04
<i>Grm3</i>	1.4E-02 ± 1.2E-03	1.4E-02 ± 4.0E-03	6.7E-03 ± 8.0E-04	7.7E-03 ± 7.3E-04	2.6E-02 ± 4.1E-03	3.6E-02 ± 7.9E-03
<i>Grm4</i>	6.8E-03 ± 3.0E-03	1.1E-02 ± 3.5E-03	1.3E-02 ± 1.0E-03	1.5E-02 ± 5.1E-03	1.6E-02 ± 3.5E-03	1.7E-02 ± 1.4E-03
<i>Grm5</i>	1.1E-01 ± 7.9E-03	2.6E-01 ± 2.2E-01	1.3E-02 ± 1.6E-03	4.9E-02 ± 1.9E-02	1.9E-02 ± 5.7E-04	2.2E-02 ± 3.0E-03
<i>Grm6</i>	2.9E-03 ± 1.3E-03	4.4E-03 ± 2.3E-03	1.0E-03 ± 5.0E-04	1.8E-03 ± 1.8E-04	nd	nd
<i>Grm7</i>	1.2E-02 ± 1.7E-03	1.5E-02 ± 3.1E-03	9.3E-03 ± 1.9E-04	1.2E-02 ± 2.0E-03	2.2E-02 ± 4.4E-03	2.8E-02 ± 2.1E-03
<i>Grm8</i>	nd	nd	nd	nd	2.5E-03 ± 6.9E-04	5.1E-03 ± 1.1E-03
<i>Slc17a6</i>	2.4E-03 ± 3.4E-04	2.4E-03 ± 7.0E-04	7.6E-04 ± 1.5E-04	<i>4.3E-04 ± 1.0E-04</i>	nd	nd
<i>Slc17a7</i>	5.9E-02 ± 6.1E-03	5.9E-02 ± 6.1E-03	5.5E-02 ± 3.4E-03	4.2E-02 ± 4.3E-03	4.4E-02 ± 3.7E-03	6.0E-02 ± 6.4E-03
<i>Slc17a8</i>	2.1E-04 ± 6.8E-05	4.1E-04 ± 1.7E-04	2.9E-04 ± 1.4E-04	4.3E-04 ± 2.5E-04	5.6E-04 ± 2.1E-04	9.4E-04 ± 2.1E-04
<i>Slc1a1</i>	3.4E-02 ± 3.2E-03	4.4E-02 ± 6.8E-03	1.5E-02 ± 3.8E-04	1.4E-02 ± 2.8E-03	3.1E-02 ± 3.8E-03	4.7E-02 ± 1.0E-02^a
<i>Slc1a6</i>	nd	nd	nd	nd	1.2E-03 ± 1.8E-04	2.1E-03 ± 5.9E-04
<i>Gls</i>	1.7E-02 ± 8.2E-04	1.1E-02 ± 1.7E-03 ^a	1.9E-02 ± 2.8E-03	1.8E-02 ± 3.2E-03	1.4E-02 ± 1.4E-03	2.1E-02 ± 2.7E-03
<i>Gls2</i>	1.6E-02 ± 3.1E-03	2.2E-02 ± 6.7E-03	7.4E-03 ± 8.0E-04	1.0E-02 ± 2.1E-03	1.1E-02 ± 5.1E-03	1.1E-02 ± 7.3E-04

nd, not detected

Data are shown as the mean ± SEM ($n = 3$) expression level relative to that of *Gapdh*.

Bold values indicate that the mean of pwSI/group > 1.5

Italicized values indicate that the mean of pwSI/group < 0.7

Values differ significantly (^a $P < 0.05$, ^b $P < 0.01$, and ^c $P < 0.001$) between group-housed and pwSI rats.

cell layers. In particular, the expression of *Grin2a* in CA1, and of *Grik4* in CA3 was markedly increased by pwSI. Seven NMDA receptor subunits are known: *Grin1*, 4 *Grin2* isoforms (*Grin2a*, *Grin2b*, *Grin2c*, and *Grin2d*), and 2 *Grin3* isoforms (*Grin3a*, *Grin3b*). The functional NMDA receptors are either di- or triheteromeric assemblies, and their physiologic properties are dependent on the subunit composition.¹⁹ In particular, the NMDA receptors that contain GRIN2A subunits play an important role in hippocampal synaptic plasticity. For example, mice lacking *Grin2a* showed attenuated long-term potentiation in CA3–CA1 and impairments in spatial learning and contextual fear conditioning.^{23,27} Unlike AMPA and NMDA receptors, which are predominantly involved with excitatory neurotransmission, the kainate receptors are thought to function mainly as modulators of excitatory

signaling.⁷ In fact, administration of the selective GRIK1 inhibitor UBP304 attenuates coordinated neuronal activity, that is, θ oscillations, in the hippocampus.¹³ Therefore, the increased expression of glutamate receptors in CA1 and CA3 pyramidal cell layers may be involved in the behavioral alterations by pwSI, particularly in hippocampus-dependent tasks.

Taken together, the present results suggest that pwSI reduced GABA_A receptor expression in the DG granule cell layer and increased the gene expression of several NMDA and kainate receptors in the CA1 and CA3 pyramidal cell layers. Although both hippocampal pyramidal cells and DG granule cells receive projections from inhibitory GABAergic interneurons,¹¹ the GABAergic inputs to DG granule cells are more important for filtering excitatory inputs from the entorhinal cortex to hippocampal circuit.⁸

Table 3. Effect of pwSI on GABAergic neuron-related gene expression in distinct hippocampal regions of F344 rats

	CA1		CA3		DG	
	Group	pwSI	Group	pwSI	Group	pwSI
<i>Gabra1</i>	1.4E-02 ± 1.7E-03	1.4E-02 ± 2.2E-03	4.3E-03 ± 4.2E-04	3.9E-03 ± 3.2E-04	1.3E-02 ± 9.6E-04	9.0E-03 ± 3.2E-04 ^c
<i>Gabra2</i>	4.4E-03 ± 8.0E-04	4.0E-03 ± 4.3E-04	3.8E-03 ± 2.2E-04	3.7E-03 ± 2.0E-04	9.6E-03 ± 9.2E-04	6.7E-03 ± 7.8E-04 ^b
<i>Gabra3</i>	nd	nd	9.7E-05 ± 1.7E-05	1.6E-04 ± 8.9E-06	3.0E-04 ± 3.0E-05	2.6E-04 ± 3.6E-05
<i>Gabra4</i>	1.1E-02 ± 2.5E-03	1.1E-02 ± 3.2E-03	3.1E-03 ± 3.3E-04	2.8E-03 ± 7.2E-05	2.6E-02 ± 1.4E-03	1.9E-02 ± 9.2E-04 ^c
<i>Gabra5</i>	4.1E-02 ± 8.9E-03	3.9E-02 ± 1.3E-02	2.7E-02 ± 3.1E-03	3.4E-02 ± 3.9E-03	3.5E-02 ± 1.1E-03	2.7E-02 ± 2.4E-03 ^a
<i>Gabra6</i>	nd	nd	nd	nd	nd	nd
<i>Gabrb1</i>	1.1E-02 ± 2.4E-03	1.2E-02 ± 2.6E-03	6.5E-03 ± 6.3E-04	8.0E-03 ± 6.1E-04	1.8E-02 ± 1.9E-04	1.5E-02 ± 1.7E-03
<i>Gabrb2</i>	1.2E-02 ± 3.0E-03	1.1E-02 ± 3.7E-03	3.0E-03 ± 1.1E-04	2.9E-03 ± 1.5E-04	1.2E-02 ± 1.4E-03	8.8E-03 ± 6.9E-04
<i>Gabrb3</i>	4.7E-03 ± 8.8E-04	5.4E-03 ± 9.7E-04	2.7E-03 ± 7.6E-05	2.6E-03 ± 3.5E-04	1.1E-02 ± 1.6E-03	7.6E-03 ± 5.9E-04 ^a
<i>Gabrg1</i>	3.1E-03 ± 8.5E-04	2.9E-03 ± 7.7E-04	1.4E-03 ± 1.5E-04	1.2E-03 ± 2.0E-04	4.4E-03 ± 6.2E-04	2.5E-03 ± 2.6E-04 ^b
<i>Gabrg2</i>	2.9E-03 ± 9.0E-04	2.3E-03 ± 6.0E-04	1.1E-03 ± 2.0E-05	1.1E-03 ± 7.4E-05	4.8E-03 ± 1.2E-03	3.1E-03 ± 2.8E-05 ^a
<i>Gabrg3</i>	2.0E-04 ± 5.1E-05	2.4E-04 ± 2.1E-05	1.2E-04 ± 3.3E-05	8.0E-05 ± 1.4E-05	5.9E-04 ± 1.7E-04	3.3E-04 ± 4.8E-05
<i>Gabrd</i>	2.6E-03 ± 5.1E-04	2.5E-03 ± 9.0E-04	4.0E-04 ± 8.7E-05	3.3E-04 ± 1.5E-05	1.9E-02 ± 2.4E-03	1.4E-02 ± 1.7E-03
<i>Gabbr1</i>	2.1E-02 ± 2.3E-03	2.9E-02 ± 6.3E-03	7.9E-03 ± 8.6E-04	1.0E-02 ± 9.8E-04	2.1E-02 ± 1.9E-03	1.8E-02 ± 2.1E-04
<i>Gabbr2</i>	2.5E-02 ± 3.9E-03	2.5E-02 ± 4.5E-03	2.2E-02 ± 2.6E-03	2.6E-02 ± 1.1E-03	4.9E-02 ± 2.6E-03	4.3E-02 ± 3.1E-03
<i>Slc32a1</i>	4.5E-04 ± 1.0E-04	4.7E-04 ± 1.4E-04	2.8E-04 ± 4.9E-06	2.3E-04 ± 1.4E-05	4.1E-04 ± 4.2E-05	6.8E-04 ± 1.2E-04
<i>Slc6a1</i>	4.1E-03 ± 1.3E-03	5.3E-03 ± 2.1E-03	2.3E-03 ± 2.9E-04	3.4E-03 ± 6.6E-04	7.5E-03 ± 1.1E-03	1.0E-02 ± 1.6E-03
<i>Gad1</i>	3.1E-03 ± 5.6E-04	2.9E-03 ± 7.2E-04	1.1E-03 ± 4.6E-05	1.1E-03 ± 1.2E-04	5.6E-03 ± 3.7E-04	6.4E-03 ± 5.2E-04
<i>Gad2</i>	2.8E-03 ± 6.5E-04	4.0E-03 ± 1.4E-03	1.2E-03 ± 9.9E-06	1.6E-03 ± 2.7E-04	5.2E-03 ± 9.4E-04	5.6E-03 ± 5.6E-04

nd, not detected

Data are shown as the mean ± SEM (*n* = 3) expression level relative to that of *Gapdh*.

Bold values indicate that the mean of pwSI/group >1.5

Italicized values indicate that the mean of pwSI/group <0.7

Values differ significantly (^a*P* < 0.05, ^b*P* < 0.01, and ^c*P* < 0.001) between group-housed and pwSI rat

Therefore, the decreased expression of GABA_A receptor genes in DG by pwSI could contribute to attenuated susceptibility to inhibitory inputs in granule cells and to enhanced excitability in downstream circuits. In addition, the expression of synthases and transporters that participate in glutamatergic neurotransmission in presynaptic cells was increased in DG, and glutamate receptor expression downstream of the circuits, that is, in CA3 and CA1, was upregulated, although the expression of GABA receptors was unaltered in both layers of pyramidal cells. In addition, these alterations in the pyramidal cell layer might contribute to enhanced hippocampal excitability. Therefore, the hippocampal circuits of pwSI rats may be in a more excitable state than that of group-housed rats.

References

1. Andersen P, Bliss TVP, Skrede KK. 1971. Lamellar organization of hippocampal excitatory pathways. *Exp Brain Res* 13:222–238.
2. Baenninger LP. 1967. Comparison of behavioural development in socially isolated and grouped rats. *Anim Behav* 15:312–323.
3. Brünig I, Scotti E, Sidler C, Fritschy JM. 2001. Intact sorting, targeting, and clustering of γ -aminobutyric acid A receptor subtypes in hippocampal neurons in vitro. *J Comp Neurol* 443:43–55.
4. Chambers MS, Atack JR, Broughton HB, Collinson N, Cook S, Dawson GR, Hobbs SC, Marshall G, Maubach KA, Pillai GV, Reeve AJ, MacLeod AM. 2003. Identification of a novel, selective GABA_A α 5 receptor inverse agonist which enhances cognition. *J Med Chem* 46:2227–2240.
5. Cheng VY, Martin LJ, Elliott EM, Kim JH, Mount HT, Taverna FA, Roder JC, Macdonald JF, Bhamri A, Collinson N, Wafford KA, Orser BA. 2006. α 5GABA_A receptors mediate the amnesic but

not sedative-hypnotic effects of the general anesthetic etomidate. *J Neurosci* 26:3713–3720.

6. Collinson N, Kuenzi FM, Jarolimek W, Maubach KA, Cothliff R, Sur C, Smith A, Otu FM, Howell O, Atack JR, McKernan RM, Seabrook GR, Dawson GR, Whiting PJ, Rosahl TW. 2002. Enhanced learning and memory and altered GABAergic synaptic transmission in mice lacking the α 5 subunit of the GABA_A receptor. *J Neurosci* 22:5572–5580.
7. Contractor A, Mülle C, Swanson GT. 2011. Kainate receptors coming of age: milestones of 2 decades of research. *Trends Neurosci* 34:154–163.
8. Coulter DA, Carlson GC. 2007. Functional regulation of the dentate gyrus by GABA-mediated inhibition. *Prog Brain Res* 163:235–243.
9. Crestani F, Keist R, Fritschy JM, Benke D, Vogt K, Prut L, Blüthmann H, Möhler H, Rudolph U. 2002. Trace fear conditioning involves hippocampal α 5 GABA_A receptors. *Proc Natl Acad Sci USA* 99:8980–8985.
10. Farrant M, Nusser Z. 2005. Variations on an inhibitory theme: phasic and tonic activation of GABA_A receptors. *Nat Rev Neurosci* 6:215–229.
11. Freund TF, Buzsáki G. 1998. Interneurons of the hippocampus. *Hippocampus* 6:347–470.
12. Goldstein PA, Elsen FP, Ying SW, Ferguson C, Homanics GE, Harrison NL. 2002. Prolongation of hippocampal miniature inhibitory postsynaptic currents in mice lacking the GABA_A receptor α 1 subunit. *J Neurol Neurophysiol* 88:3206–3217.
13. Huxter JR, Zinyuk LE, Roloff EV, Clarke VR, Dolman NP, More JC, Jane DE, Collingridge GL, Muller RU. 2007. Inhibition of kainate receptors reduces the frequency of hippocampal θ oscillations. *J Neurosci* 27:2212–2223.
14. Ibi D, Takuma K, Koike H, Mizoguchi H, Tsuritani K, Kuwahara Y, Kamei H, Nagai T, Yoneda Y, Nabeshima T, Yamada K. 2007. Social isolation rearing-induced impairment of the hippocampal neurogen-

- esis is associated with deficits in spatial memory and emotion-related behaviors in juvenile mice. *J Neurochem* **105**:921–932.
15. **Koike H, Ibi D, Mizoguchi H, Nagai T, Nitta A, Takuma K, Nabeshima T, Yoneda Y, Yamada K.** 2009. Behavioral abnormality and pharmacologic response in social isolation-reared mice. *Behav Brain Res* **202**:114–121.
 16. **Lu L, Bao G, Chen H, Xia P, Fan X, Zhang J, Pei G, Ma L.** 2003. Modification of hippocampal neurogenesis and neuroplasticity by social environments. *Exp Neurol* **183**:600–609.
 17. **Meaney MJ, Stewart J.** 1981. A descriptive study of social development in the rat (*Rattus norvegicus*). *Anim Behav* **29**:34–45.
 18. **Neves G, Cooke SF, Bliss TVP.** 2008. Synaptic plasticity, memory, and the hippocampus: a neural network approach to causality. *Nat Rev Neurosci* **9**:65–75.
 19. **Paolletti P, Bellone C, Zhou Q.** 2013. NMDA receptor subunit diversity: impact on receptor properties, synaptic plasticity, and disease. *Nat Rev Neurosci* **14**:383–400.
 20. **Parker V, Morinan A.** 1986. The socially isolated rat as a model for anxiety. *Neuropharmacology* **25**:663–664.
 21. **Quan MN, Tian YT, Xu KH, Zhang T, Yang Z.** 2010. Postweaning social isolation influences spatial cognition, prefrontal cortical synaptic plasticity, and hippocampal potassium ion channels in Wistar rats. *Neuroscience* **169**:214–222.
 22. **Sahakian BJ, Robbins TW, Iversen SD.** 1977. The effects of isolation rearing on exploration in the rat. *Anim Learn Behav* **5**:193–198.
 23. **Sakimura K, Kutsuwada T, Ito I, Manabe T, Takayama C, Kushiya E, Yagi T, Aizawa S, Inoue Y, Sugiyama H, Mishina M.** 1995. Reduced hippocampal LTP and spatial learning in mice lacking NMDA receptor $\epsilon 1$ subunit. *Nature* **373**:151–155.
 24. **Sieghart W, Sperk G.** 2002. Subunit composition, distribution and function of GABA-A receptor subtypes. *Curr Top Med Chem* **2**:795–816.
 25. **Sigel E, Steinmann ME.** 2012. Structure, function, and modulation of GABA_A receptors. *J Biol Chem* **287**:40224–40231.
 26. **Sperk G, Schwarzer C, Tsunashima K, Fuchs K, Sieghart W.** 1997. GABA_A receptor subunits in the rat hippocampus I: immunocytochemical distribution of 13 subunits. *Neuroscience* **80**:987–1000.
 27. **Sprengel R, Suchanek B, Amico C, Brusa R, Burnashev N, Rozov A, Hvalby O, Jensen V, Paulsen O, Andersen P, Kim JJ, Thompson RF, Sun W, Webster LC, Grant SG, Eilers J, Konnerth A, Li J, McNamara JO, Seeburg PH.** 1998. Importance of the intracellular domain of NR2 subunits for NMDA receptor function in vivo. *Cell* **92**:279–289.
 28. **Squire LR.** 1992. Memory and the hippocampus: a synthesis from findings with rats, monkeys, and humans. *Psychol Rev* **99**:195–231.
 29. **Syme LA.** 1973. Social isolation at weaning: some effects on 2 measures of activity. *Anim Learn Behav* **1**:161–163.
 30. **Thor DH, Holloway WR Jr.** 1984. Social play in juvenile rats: a decade of methodological and experimental research. *Neurosci Biobehav Rev* **8**:455–464.
 31. **Tretter V, Moss SJ.** 2008. GABA_A receptor dynamics and constructing GABAergic synapses. *Front Mol Neurosci* **1**:7.
 32. **Toth M, Mikics E, Tulogdi A, Aliczki M, Haller J.** 2011. Postweaning social isolation induces abnormal forms of aggression in conjunction with increased glucocorticoid and autonomic stress responses. *Horm Behav* **60**:28–36.
 33. **Vöikar V, Polus A, Vasar E, Rauvala H.** 2004. Long-term individual housing in C57BL/6J and DBA/2 mice: assessment of behavioral consequences. *Genes Brain Behav* **4**:240–252.
 34. **Wongwitdecha N, Marsden CA.** 1996. Social isolation increases aggressive behaviour and alters the effects of diazepam in the rat social interaction test. *Behav Brain Res* **75**:27–32.
 35. **Yamamuro Y.** 2008. Maternal environment alters social interactive traits but not open-field behavior in Fischer 344 rats. *Exp Anim* **57**:439–446.
 36. **Yoshioka W, Endo N, Kurashige A, Haijima A, Endo T, Shibata T, Nishiyama R, Kakeyama M, Tohyama C.** 2012. Fluorescence laser microdissection reveals a distinct pattern of gene activation in the mouse hippocampal region. *Sci Rep* **2**:1–7.