# Antidiabetic Effects and Mechanisms of Action of *Uncaria gambir* Roxb. in Diabetic Sprague–Dawley Rats

Indah Fajarwati, MS, PhD,<sup>1,\*</sup> Dedy D Solihin, Ir, DEA, PhD,<sup>2,3</sup> Tutik Wresdiyati, DVM, PhD,<sup>4</sup> Irmanida Batubara, MS, PhD,<sup>5,6</sup> and Sela S Mariya, PhD<sup>1</sup>

The antidiabetic properties of *Uncaria gambir* are not yet fully understood, particularly concerning how it affects diabetic animal models. Further investigation in this aspect is pivotal before initiating clinical evaluations. This study aimed to investigate the antidiabetic activity of *U. gambir* and how it influences blood glucose levels in diabetic Sprague–Dawley rats. In this study, 28 rats were divided into 7 groups. The groups were as follows: a nondiabetic rat group, a nondiabetic rat group given *U. gambir*, a diabetic rat group, a diabetic rat group given glibenclamide, and 3 diabetic rat groups given *U. gambir* at 3 doses (200, 300, and 400 mg/kg). Diabetes was induced using streptozotocin (50 mg/kg) given by intraperitoneal injection. Blood glucose levels were measured weekly, and the animals were euthanized at the end of the experiment. Intracardiac blood and tissues such as the pancreas, liver, and skeletal muscle were collected for further analysis. The results showed that administering *U. gambir* to diabetic rats resulted in significantly lower blood glucose levels than untreated diabetic rats. *U. gambir* has a complex mechanism to reduce blood glucose levels. including increase of insulin production, preservation of the islets and pancreatic  $\beta$  cells, and optimization of glycogenesis, as reflected in a significant increase in liver glycogen levels. These findings suggest that *U. gambir*'s multicompound and multitarget capabilities in controlling blood glucose levels may have utility for treatment of diabetes.

Abbreviations and Acronyms: DM, diabetes mellitus; Glib, glibenclamide; PAS, periodic acid-Schiff; UG, Uncaria gambir

DOI: 10.30802/AALAS-JAALAS-24-117

## Introduction

Diabetes, marked by elevated blood glucose levels or hyperglycemia,<sup>9</sup> stands as the most prevalent endocrine disorder. Globally, diabetes remains a formidable challenge for the healthcare system,<sup>41</sup> ranking among the top 10 causes of death with an estimated 6.7 million deaths in 2021.<sup>38,41</sup> Notably, it emerges as the second most common comorbidity after hypertension for COVID-19 patients in several nations, such as China, South Korea, and Indonesia.<sup>8,36</sup>

Diabetes consists of 2 main types: type 1 diabetes, the result of autoimmunity toward the pancreatic  $\beta$  cells; and type 2 diabetes, mostly linked to insufficient secretion of glucose-specific insulin.<sup>56</sup> Despite their distinct etiologies, both types share overlapping pathophysiology. For instance, both types can lead to a reduction in  $\beta$ -cell mass.<sup>13,48</sup> They are both characterized by persistent hyperglycemia, which ultimately leads to the death of  $\beta$  cells.<sup>17,37</sup> This is also causatively connected to dysfunction in pancreatic  $\beta$  cells and the decline of insulin levels.<sup>55</sup> Given the complex pathophysiology, it has been

\*Corresponding author. Email: indahfjwati@gmail.com

speculated that certain antidiabetic medications approved for type 2 diabetes treatments might confer benefits for type 1 diabetes individuals.<sup>14</sup>

Combination antihyperglycemic drugs could be of value for patients struggling to attain clinical moderation with first-line oral antidiabetic agents as monotherapy.<sup>47</sup> This complexity underscores the necessity for multifaceted approaches in diabetes treatment. In the United States, only 36% of individuals achieve glycemic control despite many medications available.<sup>47</sup> Patient nonadherence is the main contributing factor to its ineffectiveness. On the other hand, a study has reported that the efficacy of antidiabetic drugs is only 41%.<sup>1</sup> Hence, there is an urgent need for new treatment alternatives. Identifying novel antidiabetic agents with potent multimechanistic actions could offer a promising approach. Such medications would likely target multiple pathways and mechanisms involved in the development of diabetes, thereby maximizing effectiveness and potentially improving patient outcomes.

Plants have emerged as sources of herbal medicines and often serve as precursors for synthetic drugs.<sup>59</sup> Approximately half of today's medicines derive from natural sources, including plants.<sup>54</sup> Various mechanisms offered by medicinal plants contribute to lowering blood glucose levels. Several plant species exhibit a capacity to diminish blood glucose levels by inhibiting the  $\alpha$ -glycosidase enzyme.<sup>20,63</sup> Others are known to rejuvenate pancreatic  $\beta$  cells, enhance insulin secretion and sensitivity, accumulate antioxidants, impede hepatic gluconeogenesis, and alleviate metabolic syndrome in diabetic patients.<sup>16,43,53,58</sup>

Submitted: 16 Jan 2024. Revision requested: 22 Feb 2024. Accepted: 15 Aug 2024. <sup>1</sup>Center for Biomedical Research, Health Research Organization, National Research and Innovation Agency (BRIN), Cibinong, Indonesia; <sup>2</sup>Department of Biology, Faculty of Mathematics and Natural Sciences, IPB University, Dramaga, Bogor, Indonesia; <sup>3</sup>Biotech Center, IPB University, Dramaga, Bogor, Indonesia; <sup>4</sup>Divisions of Anatomy, Histology, and Embriology, School of Veterinary Medicine and Biomedical Sciences, IPB University, Dramaga, Bogor, Indonesia; <sup>5</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences, IPB University, Dramaga, Bogor, Indonesia; and <sup>6</sup>Tropical Biopharmaca Research Center, IPB University, Bogor, Indonesia

Vol 00, No 00 Journal of the American Association for Laboratory Animal Science Month 2024

Uncaria gambir, native to Southeast Asia and classified under the Rubiaceae family, is commonly known as 'Gambir' or 'Gambier' in Indonesia and Malaysia.<sup>49</sup> Locally, its leaves and twigs undergo traditional processing to produce an extract used in diverse practices, such as 'menyirih', where it is combined with other ingredients, wrapped in betel leaves, and chewed, as well as in remedies for ailments like diarrhea, dysentery, and premature ejaculation.<sup>44</sup> Numerous studies have elucidated the antidiabetic potential of *Uncaria gambir*. In vitro experiments have indicated its capability to inhibit the  $\alpha$ -glucosidase enzyme.<sup>6,39,70</sup> Furthermore, various studies have corroborated its possession of antioxidant properties.<sup>4,34</sup> Nevertheless, despite these findings, a fundamental understanding of *U. Gambir*'s antidiabetic activity in animal models of diabetes remains limited.

While many studies have used in vitro and in silico approaches, animal models are still crucial for investigating different aspects of diabetes, including its underlying causes, progression, and response to treatment.<sup>40</sup> Streptozotocin is widely used to induce experimental diabetes in animals.<sup>25</sup> It has been found to be more effective than other chemicals, like alloxan, due to its superior diabetogenic properties, and works well across different species and produces more consistent experimental outcomes.3 The diabetogenic mechanism of streptozotocin primarily revolves around the DNA methylating activity of its methyl nitrosourea component. It leads to DNA damage with resultant necrosis of the pancreatic  $\beta$  cells.<sup>24</sup> In addition, streptozotocin generates superoxide dismutase anions that act on the mitochondria and cause diabetic complications.<sup>52</sup> Given the above, the present study was conducted to investigate the potential of Uncaria gambir as an antidiabetic agent and elucidate the underlying mechanisms that contribute to its efficacy in this diabetic model.

## **Materials and Methods**

**Plant material.** Fresh leaves and young twigs of *Uncaria gambir* were obtained from a local farm in Surantih district, Pesisir Selatan regency, West Sumatera province, Indonesia. The herbarium of Andalas University in West Sumatra, Indonesia confirmed the material as being from *U. gambir*.

**Preparation of** *Uncaria gambir* **extract.** In total, 36.5 kg of fresh leaves and twigs were boiled for 2 h and then pressed using a jack press to obtain the liquid extract. Liquid extract was filtered and deposited in a wooden container called *paraku* for 24 h. In the *paraku*, the extract was allowed to cool into a paste. The extract was drained and pressed again with stone ballast to make the paste denser and immediately printable. The printed extract was then sun dried for 3 to 7 d.

**Phytochemical profile.** Phytochemical examinations were conducted using both qualitative and quantitative approaches. Following standard procedures, qualitative methods identified alkaloids, flavonoids, saponins, quinones, tannins, triterpenoids, and steroids.<sup>32</sup>

The quantitative analysis involved the measurement of catechin referring to the Indonesian National Standard (SNI 01 to 3391-2000). Briefly, catechin measurement was performed using a spectrophotometric analysis, which involved comparing the absorption of the sample solution with that of a standard solution. The standard solution, containing catechin with a purity level of over 98% and sourced from Sigma-Aldrich (St. Louis, MO), was used as a benchmark. The analysis was carried out spectrophotometrically using a wavelength of 279 nm (U-2010 spectrophotometer; Hitachi, Tokyo, Japan). Quercetin was quantitatively assessed using HPLC (HPLC-UV-VIS; Hitachi, Tokyo, Japan), following the protocols outlined in a previous study.<sup>33</sup> In summary, 1 gm of the sample was dissolved in 10 mL of MeOH tert-butylhydroquinone. The solution was sonicated for 20 min and then filtered using a 0.45-µM filter paper. The dissolved sample was then injected into the HPLC and evaluated using a wavelength of 370 nm.

Animals and housing. This investigation used 28 male Sprague-Dawley rats (Rattus norvegicus). The rats were obtained from the Indonesian Food and Drug Authority (BPOM) Jakarta, with the breeding stock originating from Charles River Laboratories (Wilmington, MA) and supplied by the Japan International Cooperation Agency (JICA). The rats were housed in a controlled environment with the temperature maintained at 22 to 24 °C, following a 12-h dark/12-h light cycle, and a relative humidity ranging from 50% to 60%. Two rats per cage were kept in nontransparent polypropylene cages, with their tails marked using a permanent marker for identification.<sup>12</sup> The size of cage was 40 cm in length, 31 cm in width, and 18 cm in height. The cages were equipped with solid floors covered with medium to large wood shavings sourced from Guazuma ulmifolia Lamk. The placement of cages on the rack was randomized to minimize the effects of environmental variables. The cage was equipped with a wire mesh top, allowing rats to observe external activities and interact with neighboring rats while ensuring optimal ventilation inside the cage. Miniplastic cups were provided for playing and gnawing in each cage for enrichment. The rats were fed commercial pellets explicitly formulated for laboratory rats, including white rats. The food had a maximum water content of 12%, a minimum protein content of 20%, a maximum fat content of 4%, a maximum crude fiber content of 4%, and a calcium content of 12% with a phosphorus content of 0.7% (Rat Bio; Citra Ina Feedmill, Jakarta, Indonesia). The water provided was bottled drinking water that met the standard quality requirements for human consumption. Food and water were provided ad libitum. The rats, aged 8 wk and weighing 230 to 250 g, were in good health and underwent a 14-d acclimation before any treatment was administered. The rats were determined to be free from internal and external parasites. Veterinary examinations were regularly conducted throughout the study period. The Committee Responsible for Animal Welfare at the Veterinary Teaching Hospital, IPB University, approved the rat handling and the study protocol under No. 29-2016 ACUC RSHP FKH-IPB. Importantly, all procedures and the utilization of animals strictly adhered to the Guide for the Care and Use of Laboratory Animals.<sup>35</sup>

**Sample size determinations.** The sample size calculation for the study was done using G\*Power software version 3.1.9.7.<sup>28</sup> The *F* test family through ANOVA (fixed effect, omnibus, one-way model, with a *P* value of 0.05 and a power of 0.8) was considered for this calculation. The chosen effect size value was 0.840, obtained from the  $\eta$  value resulting from evaluating the AUC value against blood glucose levels. A total sample size of 28 rats was determined to be required for the study, with 4 rats in each experimental group.

**Experimental design.** In this study, 28 rats were randomly divided into 7 experimental groups, each containing 4 animals. The groups were as follows: (1) Non-DM group, in which rats were administered a vehicle solution by injection and daily oral placebo administration via gavage needle; (2) DM group, in which rats received streptozotocin injection and daily oral placebo administration; (3) Non-DM + UG 400 group, where rats received a vehicle solution (sterile distilled water) by subcutaneous injection and daily oral administration of 400 mg/kg of *U. gambir*; (4) DM + Glib 2.5 group, where rats received

streptozotocin injection and daily oral administration of 2.5 mg/kg of glibenclamide; (5) DM + UG 200 group, where rats received streptozotocin injection and daily oral administration of 200 mg/kg of *U. gambir*; (6) DM + UG 300 group, where rats received streptozotocin injection and daily oral administration of 300 mg/kg of *U. gambir*; and (7) DM + UG 400 group, where rats received streptozotocin injection and daily oral given of 400 mg/kg of *U. gambir*. The dry *U. gambir* extract is dissolved in distilled water before use. The *U. gambir* solution, which tends to be bitter, was administered orally using a feeding needle. All treatments were administered daily for 28 d.

**Diabetes induction.** After an overnight fasting period, each rat was administered 50 mg/kg of streptozotocin intraperitoneally. The streptozotocin was prepared for injection by dissolving in a 0.1 M citrate buffer with a pH of 4.5. Following injection, the rats consumed a 10% sucrose solution for 24 h and were later provided with standard drinking water ad libitum.<sup>25</sup> Seven days later, fasting blood glucose levels were evaluated using samples obtained from the tail tip, and glucose was measured with a glucometer (Accu-Check Active; Roche, Mannheim, Germany) to confirm diabetes induction. Rats having a fasting blood glucose of greater than 200 mg/dL were considered to be diabetic.<sup>26</sup> Treatment of the diabetic rats commenced after 7 d of streptozotocin induction.

**Euthanasia and sample collection.** The rats were humanely euthanized by cardiac exsanguination following deep anesthesia with ketamine (80 mg/kg) and xylazine (20 mg/kg) administered intraperitoneally. The blood was collected and deposited into vacuum serum tube tubes without any added anticoagulants. Afterward, these blood samples were moved to Eppendorf tubes and centrifuged at 1,300 RCF for 10 min to separate the serum which was then collected. Immediately following euthanasia, the pancreas was collected and preserved in Bouin solution; and liver and skeltal muscle samples were dehydrated in ethanol solutions, cleared using xylene, and subsequently embedded in paraffin wax for sectioning. Three-micrometer sections of the tissues were then produced using a rotary microtome.

**Insulin level.** Serum samples were used to determine the insulin level of rats. Insulin levels were assessed using The Rat Insulin ELISA Kit (catalog no. 90010, Crystal Chem, Inc., Elk Grove Village, IL). The assay was conducted in adherence to the manufacturer's guidelines and protocols.

Immunohistochemistry staining. After dewaxing and rehydrating, the tissue sections of the pancreas were incubated in 0.03% hydrogen peroxide, washed using 10% PBS, and incubated in 5% normal serum for 50 min and background sniper for 15 min. Subsequently, the samples were incubated with rat anti-insulin antibody (Genetex GTX28304, Genetex, Inc., Irvine, CA) (diluted in PBS 1:500) for 1 d. The samples underwent 3 washes in a PBS solution, followed by a 20-min incubation with Trekkie Universal Link (Biocare Medical, Inc., Pacheco, CA). Subsequently, the samples were rinsed again in PBS and incubated for 20 min with Trek-Avidin HRP (Biocare Medical). The sections were visualized with DAB (40%), rinsed, stained with hematoxylin, dehydrated, cleared, and well coverslipped. The tissues were examined under light microscopy using an Olympus CX31 Microscope with Indomicro Camera (Indomicro, Jakarta, Indonesia). In total, 15 visual fields per group were randomly chosen for evaluation.

**Periodic acid–Schiff staining.** Periodic acid–Schiff (PAS) staining was employed to visualize liver and skeletal muscle glycogen deposits. Paraffin sections of the organs were

deparaffinized and incubated in 0.5% periodic acid (Sigma, Deisenhofen, Germany) for 5 min. Following these steps, the sections were rinsed in distilled water for 5 min, immersed in Schiff reagent (Sigma-Aldrich, St. Louis, MO) for 15 min, and then subjected to another 5-min wash in distilled water. Subsequently, they were counterstained with hematoxylin for 1 min, rinsed again in distilled water for 5 min, dehydrated, and finally mounted for observation. The staining results were observed and documented using a light microscope equipped with a camera and analyzed using the ImageJ application. Analysis was performed on 30 visual fields per group.

**Statistical analysis.** Before further analysis, testing the data for normality and homogeneity was crucial. If the data adhered to those criteria, the ANOVA test was initially performed, and the Duncan multiple range test was applied to explore and compare the variations among the groups in the study. Specifically for the AUC of glucose values, a post hoc test was performed using the Dunnett test, with the DM group as the control. In cases where the dataset deviates from normality and homogeneity, the Kruskal-Wallis test is used, followed by the Mann-Whitney test. Significant differences were indicated as  $P \le 0.05$ . The outcomes are depicted as mean ± SEM. Statistical analyses were conducted using version 21.0 of SPSS (Chicago, IL).

## Results

**Phytochemical profile of** *Uncaria gambir* **extract.** The analysis of the bioactive aqueous extract of *Uncaria gambir* indicated the presence of flavonoids, tannins, saponins, and triterpenoids while absent for alkaloids, quinones, and steroids. The HPLC analysis revealed the presence of querce-tin, as shown in the chromatogram (Figure 1), indicating 2.65 mg/g of quercetin. In addition, as the marker for assessing the quality of the *U. gambir* extracts, the total catechin content was  $80.56\% \pm 0.57$ , which agreed with National Standard Indonesia (SNI) No. 01 to 3391-2000 and was categorized as grade 1 in the trade market.

The impact of Uncaria gambir extract on blood glucose level. Blood glucose levels in diabetic rats treated with *U. gambir* at a dose of 200 mg/kg body weight returned to normal from severe (Figure 2A). By day 16, this group's average blood glucose level had decreased to 105 mg/dL from an initial level of 338 mg/ dL. In contrast, rats in other treatment groups did not demonstrate similar reductions in blood glucose. The administration of *U. gambir* at doses of 300 and 400 mg/kg BW, as well as administration of glibenclamide, only managed to lower blood glucose levels to the intermediate level.

The influence of *U. gambir* on fluctuations in blood glucose levels over 28 d was further statistically evaluated by analyzing AUC (Figure 2B). The results showed that diabetic rats treated with 200 mg/kg of *U. gambir* had significantly lower blood glucose levels than untreated diabetic rats (DM group). This result suggests that administration 200 mg/kg of *U. gambir* to diabetic rats could reduce high blood glucose levels. However, diabetic rats treated with higher doses of *U. gambir* (300 and 400 mg/kg) and those treated with glibenclamide did not show significant differences in blood glucose levels compared with those of untreated diabetic rats.

**The effect of** *Uncaria gambir* **extract on insulin level.** The insulin levels were significantly lower in diabetic rats (DM) than in nondiabetic rats (Non-DM). After being treated with *U. gambir*, diabetic rats exhibited significantly higher insulin levels when compared with the untreated diabetic rat group. This increase was observed in all groups of rats treated with different doses



Figure 1. HPLC-UV-VIS quercetin profile of *Uncaria gambir*. (A) Quercetin in *Uncaria gambir* extract. (B) Quercetin standard. The red letter 'q' at the peak of the chromatogram indicates quercetin.

of *U. gambir* (DM + UG 200, DM + UG 300, and DM + UG 400). Furthermore, the insulin levels in these groups (DM + UG 200, DM + UG 300, and DM + UG 400) were not significantly different from nondiabetic rats (Non-DM). Conversely, diabetic rats treated with glibenclamide (DM + Glib 2.5) displayed no significant improvement, as their serum insulin levels were akin to those in the untreated diabetic rat group (DM) (Figure 3).

The effect of Uncaria gambir on the Langerhans β cells. Insulinpositive pancreatic  $\beta$  cells were cells were identified as those that stained a reddish-brown color (Figure 4A). In the diabetic rat group (DM), the β-cell number was reduced deficient. In contrast, those diabetic rats given *U. gambir* (DM + UG 200, DM + UG 300, and DM + UG 400) had more  $\beta$  cells than those of diabetic rats (DM). This result demonstrates that administration of *U. gambir* to diabetic rats could sustain the  $\beta$  cell population, although the quantity of  $\beta$  cells observed was not as extensive as those found in nondiabetic rats.

The diabetic rat groups receiving *U. gambir* extract (DM + UG 200, DM + UG 300, and DM + UG 400) quantitatively exhibited significantly higher  $\beta$ -cell percentages than the untreated

diabetic rat group (DM), although lower than the  $\beta$ -cell percentage observed in the nondiabetic group (Non-DM) (Figure 4B). Diabetic rats treated with glibenclamide (DM + Glib 2.5) did not display a notably higher  $\beta$ -cell percentage than the untreated diabetic rat group (DM). Furthermore, administration of *U. gambir* extract at 400 mg/kg to nondiabetic rats (Non-DM + G 400) did not impact the  $\beta$ -cell percentage, as evidenced by the equivalent percentage of  $\beta$  cells between this group (Non-DM + UG 400) and the nondiabetic rat group (Non-DM).

**The effect of** *Uncaria gambir* **on Langerhans islets.** The diabetic rats (DM) had the lowest count of islets compared with all other groups (Figure 4C). However, when diabetic rats were given *U. gambir* at any dosage (DM + UG 200, DM + UG 300, and DM + UG 400), they had more islets than untreated diabetic rats. Even though improvement was evident in the diabetic rats that underwent treatment, the number of islets remained below that observed in the nondiabetic rat group (Non-DM). In addition, administration of glibenclamide to diabetic rats (DM + Glib 2.5) did not significantly affect the number of islets in diabetic rats.



**Figure 2.** The effect of *Uncaria gambir* on blood glucose level. (A) Fluctuations in blood glucose levels during 28 d of observation. (B) AUC of glucose level. \*Significant difference compared with the DM group according to the Dunnett test ( $P \le 0.05$ ). DM, diabetes mellitus (DM); Glib, glibenclamide; UG, *Uncaria gambir*.



**Figure 3.** The effect of *Uncaria gambir* to insulin level. Different letters above the bar indicate significant differences according to the Duncan multiple range test ( $P \le 0.05$ ). DM, diabetes mellitus; Glib, glibencla-mide; UG, *Uncaria gambir*.

**The effect of** *Uncaria gambir* **on liver glycogen.** Staining of liver samples with PAS demonstrated (Figure 5A). The magenta color staining was more widespread in the group of nondiabetic rats (Non-DM) than in the group of diabetic rats (DM) due to a positive PAS reaction. This result was also observed in the groups that received *U. gambir* at any dosage (DM + UG 200, DM + UG 300, and DM + UG 400) and the diabetic rat group given glibenclamide (DM + Glib 2.5) stained more intensely for glycogen compared with the diabetic group (DM). PAS staining was even distributed across the liver in the nondiabetic rat group given

*U. gambir* 400 mg/kg (Non-DM + UG 400). Qualitatively similar results show that the diabetic rats had the lowest glycogen levels compared with other groups, while the treated diabetic groups demonstrated higher levels (Figure 5B).

The effect of *Uncaria gambir* on skeletal muscle glycogen. It was observed that the skeletal muscle cells of diabetic rats (DM) had less intense PAS staining for glycogen than the other groups (Figure 6A). However, there were no significant statistical differences between the groups (Figure 6B).

### Discussion

This study aimed to determine whether *Uncaria gambir* could act as an antidiabetic agent in a rat model of diabetes. Our hypothesis is that *U. gambir* may exert antidiabetic effects via a multitarget mechanism of action. Prior studies have demonstrated the significant potential of the various components of *U. gambir*.<sup>5,7,39,45,49</sup> These findings suggest that *U. gambir* may have multiple target actions in regulating blood glucose levels, and its constituents may work synergistically.

A rat model of diabetes was used in this study and produced by administration of streptozotocin at 50 mg/kg. The rats exhibited various pathophysiologies associated with diabetes, such as persistent hyperglycemia and diminished insulin levels, islets,  $\beta$  cells, and liver glycogen content. Intriguingly, treatment with *U. gambir* improved some of these pathophysiologies, thus supporting the hypothesis that *U. gambir* acts through various mechanisms to decrease blood glucose levels.



**Figure 4.** The effect of *Uncaria gambir* on islets of Langerhans and  $\beta$  cells. (A) Photomicrograph of immunohistochemical staining of insulin-activated pancreatic  $\beta$  cells; scale = 50 µm; 10 × 40 magnifications. (B) The proportion of  $\beta$  cells to the total number of cells. (C) The number of islets per high-powered field; 10 × 10 magnifications. Different letters above the bar indicate significant differences according to the Duncan multiple range test for the proportion of  $\beta$  cells or Mann-Whitney test for the number of islets ( $P \le 0.05$ ). Arrows are pancreatic  $\beta$ -cell representation. DM, diabetes mellitus; Glib, glibenclamide; L, islet of Langerhans; UG, *Uncaria gambir*.

Vol 00, No 00 Journal of the American Association for Laboratory Animal Science Month 2024



**Figure 5.** The effect of *Uncaria gambir* on liver glycogen. (A) Photomicrograph of PAS staining of liver glycogen; scale = 50  $\mu$ m. (B) Quantitative liver glycogen intensity per high-powered field; 10 × 40 magnifications. Different letters above the bar indicate significant differences according to the Duncan multiple range test (*P* ≤ 0.05). CV, central vein; DM, diabetes mellitus; Glib, glibenclamide; UG, *Uncaria gambir*.

Our results show that administration of *U. gambir* is associated with lower blood glucose levels in a diabetic rat model. The most effective dosage of *U. gambir* is 200 mg/kg, and we showed that this dosage can return blood glucose back to normal levels. According to a previous study, Sprague-Dawley rats have a fasting normal blood glucose level below 151 mg/dL.<sup>26</sup> In this study, diabetic rats treated with *U. gambir* at a dosage of 200 mg/kg achieved normal blood glucose levels from week 12 to week 20, with the best average blood glucose level (105 mg/dL) reached at week 16. Interestingly, extending the administration time and U. gambir dosage did not result in further reductions of blood glucose levels. This finding may be related to U-shaped dose-response relationships (often termed hormesis). Hormesis is a phenomenon where a substance exhibits a dual response, stimulating beneficial effects at low doses or concentrations and causing inhibitory or detrimental effects at higher doses or concentrations.<sup>15,61</sup> Hormesis has been documented in numerous biologic, toxicological, and pharmacological agents, including the effects of metformin as an antidiabetic drug,<sup>51</sup> as well as the influence of alcohol on blood sugar levels.<sup>57</sup> Perhaps, a hormetic response affects U. gambir's ability to moderate blood glucose levels, with efficacy demonstrated at one dosage but not at greater dosages. However, further research is required to confirm this effect and elucidate the underlying mechanisms.

When administered to diabetic rats, *U. gambir* increased insulin secretion. Remarkably, the diabetic rats treated with

U. gambir had insulin levels similar to those of nondiabetic rats, even though the number of Langerhans islets and  $\beta$  cells between the 2 groups was significantly different. The ability of U. gambir to increase insulin secretion could be attributed to the quercetin and catechins component of the preparation. The phytochemical test results indicate that the gambir used in the study contained a significant amount of catechin, up to 80%, while the quercetin content is measured at 2.65 mg/g. Quercetin could enhance insulin release by increasing Ca<sup>2+</sup> uptake.<sup>10</sup> Catechin has several mechanisms to regulate diabetes. Indeed, a significant aspect involves enhancing mitochondrial function by maintaining oxidative phosphorylation and ATP production, encouraging mitochondrial biogenesis, and facilitating cellular protection within mitochondria.<sup>64</sup> Mitochondria play a vital role in cell viability, and the insulin secretion triggered by glucose is contingent upon mitochondrial respiration through the electron transport chain. In addition, epicatechin could act as a signaling molecule to stimulate insulin secretion by activating calmodulin-dependent protein kinase II (CaMKII).<sup>67</sup> These available studies provide evidence supporting the role of catechin and quercetin in increasing insulin secretion. However, this does not eliminate the possibility of other phytochemicals contributing to this action. As a result, further exploration is essential to determine the involvement of other phytochemicals in insulin secretion.

In this study, we show that *U. gambir* can help to maintain the number of Langerhans' islets and  $\beta$  cells in diabetic animals.



**Figure 6.** The effect of *Uncaria gambir* on skeletal muscle glycogen. (A) Photomicrograph of PAS staining of skeletal muscle glycogen; scale = 50 µm. (B) Quantitative skeletal muscle glycogen intensity. DM, diabetes mellitus; Glib, glibenclamide; ns, not significant; UG, *Uncaria gambir*.

Even though the group of diabetic rats given *U. gambir* had fewer Langerhans islets and  $\beta$  cells than the nondiabetic rat group, the number of islets and cells was still significantly greater than that in the diabetic rat group. Langerhans islets regulate glucose homeostasis and are vital components of the pancreas.<sup>27</sup> The number of Langerhans islets and pancreatic  $\beta$  cells could be maintained through protection, regeneration, or both. Antioxidant compounds could facilitate protection, preventing  $\beta$  cells from being damaged.<sup>23</sup> Beta cells have lower antioxidant defenses than others, making them more vulnerable to oxidative damage.<sup>46</sup> Beta cells have fewer prosurvival Bcl2 proteins, making them more susceptible to damage than others.<sup>60</sup>

U. gambir has been identified as a potential source of antioxidant activity.<sup>34</sup> Catechin, an abundant component of U. gambir, is a powerful antioxidant, and the Indonesian National Standard (SNI) has designated catechin as a biomarker compound for evaluating U. gambir quality. Others have suggested that catechin could significantly enhance the activity of antioxidant enzymes, lower the levels of reduced glutathione, and decrease malondialdehyde content in the livers and kidneys of streptozotocin-induced diabetic rats.<sup>50</sup> The high antioxidant activity in *U. gambir* could contribute to its antidiabetic effects, exceeding those of glibenclamide. Unlike glibenclamide, which does not contain antioxidants, antioxidants in *U. gambir* could protect  $\beta$  cells from damage. Several studies have reported that glibenclamide controls blood glucose levels by stimulating insulin secretion.<sup>29</sup> In contrast, another study suggested that glibenclamide administration can inhibit insulin secretion as part of its mechanism for repairing pancreatic  $\beta$  cells.<sup>71</sup>

Apart from protection, the number of Langerhans islets could also be maintained through regeneration. Recent studies have found that the pancreas has a significant self-renewal capacity.<sup>27</sup> Several pancreatic regeneration mechanisms have been proposed, including neogenesis of Langerhans islets through the proliferation and differentiation of duct cells,<sup>62</sup> endogenous progenitor cells,<sup>66</sup> preexisting  $\beta$  cells,<sup>21</sup> transdifferentiation of other endocrine cells,<sup>11</sup> and self-duplication.<sup>22</sup> Previous studies have reported that  $\beta$ -cell regeneration occurs 1 to 2 wk after pancreas ligation in adult *Mus musculus*.<sup>62</sup> Similarly, newly formed small Langerhans islets were found one week after pancreatic ligation in *Rattus norvegicus*.<sup>62</sup>

Various plants have also been reported to enhance β-cell function and regeneration, including Centaurium erythraea, Cornus officinalis, Lactarius deterrimus, Panax ginseng, and Gardenia jasminoides.65 Saponin and quercetin in plants have specific effects on  $\beta$  cells that may be useful in treating diabetes.<sup>65</sup> Quercetin not only enhances insulin secretion but also prompts the regeneration of pancreatic  $\beta$  cells.<sup>31</sup> Saponins could facilitate  $\beta$ -cell regeneration by protecting cells from apoptosis and promoting cell proliferation, thereby safeguarding Langerhans islets from damage.<sup>18</sup> Saponins activate the Wnt/β-catenin/ TCF7L2 pathway, initiating the release of GLP-1 and GIP, thereby fostering the proliferation of  $\beta$  cells.<sup>18</sup> The Wnt signaling pathway is pivotal in developing the pancreatic endocrine system and regulating β-cell functions such as insulin secretion, cell survival, and proliferation.<sup>18</sup> TCF7L2 influences the growth and differentiation of cells in Langerhans islets.<sup>18</sup> One study<sup>68</sup> suggested that saponin derived from Astragalus membranaceus notably boosted the expression of genes related to β-cell differentiation and also elevate the mRNA levels of Neurogenin 3 (Ngn3), recognized as a marker for endocrine progenitors. Hence, this study implies that U. gambir could potentially stimulate the regeneration of Langerhans islets and

pancreatic  $\beta$  cells. Further investigation is required to validate the molecular mechanisms through which *U. gambir* sustains pancreatic  $\beta$  cells.

*U. Gambir's* antidiabetic properties could be attributed to its ability to maintain liver glycogen content, which helps to reduce glucose output in the bloodstream. The present study found that diabetic rats had significantly lower glycogen content in hepatocytes compared with nondiabetic rats. Meanwhile, rats treated with *U. gambir* at varying dosages (200, 300, and 400 mg/kg) had higher glycogen content than diabetic rats, as did rats given glibenclamide. These findings indicate that *U. gambir* could effectively increase liver glycogen content in diabetic rats. The liver regulates blood glucose levels by balancing glycolysis, gluconeogenesis, and glycogen metabolism.<sup>2</sup> Therefore, it is one of the primary targets for diabetes treatment.<sup>30</sup> Inhibiting hepatic glucose release (gluconeogenesis) could be achieved by increasing glycogen storage.<sup>2</sup>

Metformin is an antidiabetic drug that also works in the liver by inhibiting gluconeogenesis.<sup>42</sup> The hepatic glycogen level can be a biomarker for evaluating the antihyperglycemic effects of various drugs.<sup>69</sup> Catechin is one of the phytochemicals in *U. gambir* that is believed to be responsible for maintaining hepatic glycogen content. Catechin increases glucokinase, glucose-6-phosphate, glycogen synthase, and glycogen phosphorylase to normal levels in diabetic rats.<sup>19</sup> Catechin also enhances the expression of GLUT4 protein and mRNA.<sup>19</sup>

This study also attempted to investigate the mechanism of *U. gambir* on glycogen content in skeletal muscles. Regrettably, the diabetes model used in this study was not accompanied by a decrease in skeletal muscle glycogen levels, posing a challenge to identifying the effect of *U. Gambir* on skeletal muscle glycogen.

Our results offer compelling evidence supporting the potential of *U. gambir* as a highly effective treatment for various pathophysiologies linked to diabetes. The results suggest that *U. gambir* holds promise in alleviating a spectrum of diabetes-related conditions, positioning it as a versatile and promising antidiabetic agent. By targeting multiple pathophysiologies simultaneously, *U. gambir* could provide an efficient and holistic approach to managing diabetes. Overall, these findings highlight the need for further research into the potential benefits of *U. gambir* for diabetes management.

## Acknowledgments

Not applicable.

### **Conflict of Interest**

The authors have no conflicts of interest to declare.

#### Funding

This work was internally funded.

## **Author Contributions**

Indah Fajarwati: conceptualization, methodology, investigation, resources, formal analysis, writing. Dedy Duryadi Solihin: conceptualization, supervision, resources, validation. Tutik Wresdiyati: conceptualization, supervision, resources, methodology. Irmanida Batubara: conceptualization, supervision, validation. Sela S Mariya: writing- reviewing and editing, formal analysis, visualization.

## Data Sharing Statement

Most of the data are included in the manuscript. Additional can be found from the corresponding author based on reasonable request.

#### References

- 1. Agarwal A, Jadhav P, Deshmukh Y. 2014. Prescribing pattern and efficacy of anti-diabetic drugs in maintaining optimal glycemic levels in diabetic patients. J Basic Clin Pharm 5:79–83.
- Agius L. 2007. New hepatic targets for glycaemic control in diabetes. Best Pract Res Clin Endocrinol Metab 21:587–605.
- 3. Alema NM, Periasamy G, Sibhat GG, Tekulu GH, Hiben MG. 2020. Antidiabetic activity of extracts of *Terminalia brownii* fresen. Stem bark in mice. J Exp Pharmacol **12**:61–71.
- Amir M, Mujeeb M, Khan A, Ashraf K, Sharma D, Aqil M. 2012. Phytochemical analysis and in vitro antioxidant activity of *Uncaria* gambir. Int J Green Pharm 6:67–72.
- Arundita S, Ismed F, Rita RS, Putra DP. 2020. (+)-Catechin & proanthocyanidin fraction of *Uncaria gambir* Roxb. Improve adipocytes differentiation & glucose uptake of 3t3-11 cells via sirtuin-1, peroxisome proliferator-activated receptor γ (PPAR γ), glucose transporter type 4 (GLUT-4) expressions. Adv Pharm Bull 10:602–609.
- Arundita S, Kurniawan F, Ismed F, Rita RS, Putra DP. 2020. In vitro alpha glucosidase activity of *Uncaria gambir* Roxb. and syzygium polyanthum (wight) walp. From West Sumatra, Indonesia. Open Access Maced J Med Sci 8:810–817.
- Azeem M, Hanif M, Mahmood K, Ameer N, Chughtai FRS, Abid U. 2023. An insight into anticancer, antioxidant, antimicrobial, antidiabetic and anti-inflammatory effects of quercetin: A review. Polym Bull (Berl) 80:241–262.
- Bajgain KT, Badal S, Bajgain BB, Santana MJ. 2021. Prevalence of comorbidities among individuals with COVID-19: A rapid review of current literature. Am J Infect Control 49:238–246.
- Balwan WK, Saba N, Zargar JI. 2022. Burden of diabetes and role of medicinal plants in its treatment. Saudi J Med Pharm Sci 8:355–361.
- Bardy G, Virsolvy A, Quignard JF, Ravier MA, Bertrand G, Dalle S, Cros G, Magous R, Richard S, Oiry C. 2013. Quercetin induces insulin secretion by direct activation of L-type calcium channels in pancreatic beta cells. Br J Pharmacol 169:1102–1113.
- Ben-Othman N, Vieira A, Courtney M, Record F, Gjernes E, Avolio F, Hadzic B, et al. 2017. Long-term GABA administration induces alpha cell-mediated beta-like cell neogenesis. Cell 168:73–85.e11.
- 12. Burn CC, Mazlan NHB, Chancellor N, Wells DJ. 2021. The pen is milder than the blade: Identification marking mice using ink on the tail appears more humane than ear-punching even with local anaesthetic. Animals 11:1644.
- Butler AE, Janson J, Bonner-Weir S, Ritzel R, Rizza RA, Butler PC. 2003. Beta-cell deficit and increased beta-cell apoptosis in humans with type 2 diabetes. Diabetes 52:102–110.
- Cai X, Lin C, Yang W, Nie L, Ji L. 2021. Non-insulin antidiabetes treatment in type 1 diabetes mellitus: A systematic review and meta-analysis. Diabetes Metab J 45:312–325.
- Calabrese EJ, Baldwin LA. 2001. U-shaped dose-responses in biology, toxicology, and public health. Annu Rev Public Health 22:15–33.
- Choudhury H, Pandey M, Hua CK, Mun CS, Jing JK, Kong L, Ern LY, et al. 2018. An update on natural compounds in the remedy of diabetes mellitus: A systematic review. J Tradit Complement Med 8:361–376.
- 17. **Costes S, Bertrand G, Ravier MA.** 2021. Mechanisms of beta-cell apoptosis in type 2 diabetes-prone situations and potential protection by GLP-1-based therapies. Int J Mol Sci **22**:5303.
- Cui J, Duan J, Chu J, Guo C, Xi M, Li Y, Weng Y, et al. 2020. Chikusetsu saponin IVa protects pancreatic β cell against intermittent high glucose-induced injury by activating Wnt/β-catenin/TCF7L2 pathway. Aging (Albany NY) 12:1591–1609.
- Daisy P, Balasubramanian K, Rajalakshmi M, Eliza J, Selvaraj J. 2010. Insulin mimetic impact of catechin isolated from Cassia fistula on the glucose oxidation and molecular mechanisms of glucose uptake on streptozotocin-induced diabetic Wistar rats. Phytomedicine 17:28–36.
- Dirir AM, Daou M, Yousef AF, Yousef LF. 2022. A review of alpha-glucosidase inhibitors from plants as potential candidates for the treatment of type-2 diabetes. Phytochem Rev 21: 1049–1079.

- 21. **Dong S, Wu H.** 2018. Regenerating  $\beta$  cells of the pancreas—Potential developments in diabetes treatment. Expert Opin Biol Ther **18**:175–185.
- Dor Y, Brown J, Martinez OI, Melton DA. 2004. Adult pancreatic b-cells are formed by self-duplication rather than stem-cell differentiation. Nature 429: 41–46.
- Eguchi N, Vaziri ND, Dafoe DC, Ichii H. 2021. The role of oxidative stress in pancreatic β cell dysfunction in diabetes. Int J Mol Sci 22:1–18.
- 24. Eleazu CO, Eleazu KC, Chukwuma S, Essien UN. 2013. Review of the mechanism of cell death resulting from streptozotocin challenge in experimental animals, its practical use and potential risk to humans. J Diabetes Metab Disord **12**:60.
- 25. Fajarwati I, Solihin DD, Wresdiyati T, Batubara I. 2023. Administration of alloxan and streptozotocin in Sprague Dawley rats and the challenges in producing diabetes model. IOP Conf Ser Earth Environ Sci 1174:012035.
- 26. Fajarwati I, Solihin DD, Wresdiyati T, Batubara I. 2023. Self-recovery in diabetic Sprague Dawley rats induced by intraperitoneal alloxan and streptozotocin. Heliyon 9:e15533.
- 27. Falcomatà C, Saur D. 2021. Self-renewal equality in pancreas homeostasis, regeneration, and cancer. Cell Rep 37:110135–3.
- Faul F, Erdfelder E, Lang A-G, Buchner A. 2007. G\* Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. Behav Res Methods 39:175–191.
- 29. Fuhlendorff J, Rorsman P, Kofod H, Brand CL, Rolin B, Mackay P, Shymko R, Carr RD. 1998. Stimulation of insulin release by repaglinide and glibenclamide involves both common and distinct processes. Diabetes 47:345–351.
- 30. Gastaldelli A, Stefan N, Häring HU. 2021. Liver-targeting drugs and their effect on blood glucose and hepatic lipids. Diabetologia 64:1461–1479.
- 31. Gurav M, Bhise S, Warghade S. 2018. Effect of quercetin on beta cell regeneration. Asian J Pharm Pharmacol 4:214–221.
- 32. **Harborne JB.** 1973. Methods of plant analysis, 3rd ed. London (UK): International Thomson Publishing.
- Hertog MGL, Hollman PCH, Venema DP. 1992. Optimization of a quantitative HPLC determination of potentially anticarcinogenic flavonoids in vegetables and fruits. J Agric Food Chem 40:1591–1598.
- 34. Hidayati MD, Rahmatulloh A. 2022. Antioxidant activity of Uncaria gambir (Hunter) Roxb extracts. Trop J Nat Prod Res 6:1215–1218. https://doi.org/10.26538/tjnpr/v6i8.9.
- 35. Institute for Laboratory Animal Research, National Research Council., Guide for the care and use of laboratory animals, 8th ed, Washington (DC): National Academies Press., 2011, https://urldefense.com/ v3/\_https:/grants.nih.gov/grants/olaw/guide-for-the-care-anduse-of-laboratory-animals.pdf.
- Karyono DR, Wicaksana AL. 2020. Current prevalence, characteristics, and comorbidities of patients with COVID-19 in Indonesia. J Community Empowerment for Health 3:77.
- 37. Khin PP, Lee JH, Jun HS. 2023. Pancreatic beta-cell dysfunction in type 2 diabetes. Eur J Inflamm 21:1721727X2311541-13.
- 38. Kim D, Alshuwaykh O, Sandhu KK, Dennis BB, Cholankeril G, Ahmed A. 2022. Trends in all-cause and cause-specific mortality among individuals with diabetes before and during the COVID-19 pandemic in the U.S. Diabetes Care 45:e107–e109.
- Kim TH. 2016. A novel α-glucosidase inhibitory constituent from Uncaria gambir. J Nat Med 70:811–815.
- 40. Kottaisamy CPD, Raj DS, Prasanth Kumar V, Sankaran U. 2021. Experimental animal models for diabetes and its related complications—A review. Lab Anim Res **37**:23.
- 41. Lin X, Xu Y, Pan X, Xu J, Ding Y, Sun X, Song X, Ren Y, Shan PF. 2020. Global, regional, and national burden and trend of diabetes in 195 countries and territories: An analysis from 1990 to 2025. Sci Rep **10**:14790.
- Liu Y, Wang C, Li J, Mei Y, Liang Y. 2019. Hypoglycemic and hypolipidemic effects of phellinus linteus mycelial extract from solid-state culture in a rat model of type 2 diabetes. Nutrients 11:296.
- 43. Luo K, Huang W, Qiao L, Zhang X, Yan D, Ning Z, Ma C, et al. 2022. *Dendrocalamus latiflorus* and its component rutin exhibit glucose-lowering activities by inhibiting hepatic glucose production via AKT activation. Acta Pharm Sin B **12**:2239–2251.

- Mat Saad MF, Goh HH, Rajikan R, Tuan Yusof TR, Baharum SN, Bunawan H. 2020. From phytochemical composition to pharmacological importance. Trop J Pharm Res 19:1767–1773.
- 45. Mechchate H, Es-Safi I, Haddad H, Bekkari H, Grafov A, Bousta D. 2021. Combination of Catechin, Epicatechin, and Rutin: Optimization of a novel complete antidiabetic formulation using a mixture design approach. J Nutr Biochem 88:108520.
- 46. Miki A, Ricordi C, Sakuma Y, Yamamoto T, Misawa R, Mita A, Molano RD, Vaziri ND, Pileggi A, Ichii H. 2018. Divergent antioxidant capacity of human islet cell subsets: A potential cause of beta-cell vulnerability in diabetes and islet transplantation. PLoS One 13:e0196570–16.
- Miller BR, Nguyen H, Hu CJH, Lin C, Nguyen QT. 2014. New and emerging drugs and targets for type 2 diabetes: Reviewing the evidence. Am Health Drug Benefits 7:452–461.
- Mukherjee N, Lin L, Contreras CJ, Templin AT. 2021. β-cell death in diabetes: Past discoveries, present understanding, and potential future advances. Metabolites 11:796.
- 49. Munggari IP, Kurnia D, Deawati Y, Julaeha E. 2022. Current research of phytochemical, medicinal and non-medicinal uses of *Uncaria gambir* Roxb.: A review. Molecules **27**:1–6.
- 50. Nazir N, Zahoor M, Ullah R, Ezzeldin E, Mostafa GAE. 2021. Curative effect of catechin isolated from *Elaeagnus umbellata* Thunb. berries for diabetes and related complications in streptozotocin-induced diabetic rats model. Molecules **26**:19.
- Panfoli I, Puddu A, Bertola N, Ravera S, Maggi D. 2021. The hormetic effect of metformin: "less is more"? Int J Mol Sci 22: 6297.
- Papaccio G, Pisanti FA, Latronico MVG, Ammendola E, Galdieri M. 2000. Multiple low-dose and single high-dose treatments with streptozotocin do not generate nitric oxide. J Cell Biochem 77:82–91.
- Pereira ASP, Banegas-Luna AJ, Peña-García J, Pérez-Sánchez H, Apostolides Z. 2019. Evaluation of the anti-diabetic activity of some common herbs and spices: Providing new insights with inverse virtual screening. Molecules 24:4030.
- 54. Qi L-W, Liu E-H, Chu C, Peng Y-B, Cai H-X, Li P. 2010. Anti-diabetic agents from natural products—An update from 2004 to 2009. Curr Top Med Chem 10:434–457.
- 55. **Shyr ZA, Wang Z, York NW, Nichols CG, Remedi MS.** 2019. The role of membrane excitability in pancreatic β-cell glucotoxicity. Sci Rep **9**:6952.
- Skyler JS, Bakris GL, Bonifacio E, Darsow T, Eckel RH, Groop L, Groop P-H, et al. 2017. Differentiation of diabetes by pathophysiology, natural history, and prognosis. Diabetes 66:241–255.
- Steiner JL, Crowell KT, Lang CH. 2015. Impact of alcohol on glycemic control and insulin action. Biomolecules 5:2223–2246.

- 58. Syukri Y, Taher M, Martien R, Lukitaningsih E, Nugroho AE, Zakaria ZA. 2021. Self-nanoemulsifying delivery of andrographolide: Ameliorating islet beta cells and inhibiting adipocyte differentiation. Adv Pharm Bull 11:171–180.
- Vuorela P, Leinonen M, Saikku P, Tammela P, Rauha J, Wennberg T, Vuorela H. 2004. Natural products in the process of finding new drug candidates. Curr Med Chem 11:1375–1389.
- 60. Wali JA, Thomas HE. 2015. Pancreatic alpha cells hold the key to survival. EBioMedicine 2:368–369.
- Wan Y, Liu J, Mai Y, Hong Y, Jia Z, Tian G, Liu Y, Liang H, Liu J. 2024. Current advances and future trends of hormesis in disease. NPJ Aging 10:26.
- 62. Wang RN, Kliippel G, Bouwens L, Jette C. 1995. Duct- to islet-cell differentiation and islet growth in the pancreas of duct-ligated adult rats. Diabetologia **38**:1405–1411.
- 63. Wang X, Liu Q, Zhu H, Wang H, Kang J, Shen Z, Chen R. 2017. Flavanols from the *Camellia sinensis* var. *assamica* and their hypoglycemic and hypolipidemic activities. Acta Pharm Sin B 7:342–346.
- 64. Wen L, Wu D, Tan X, Zhong M, Xing J, Li W, Li D, Cao F. 2022. The role of catechins in regulating diabetes: An update review. Nutrients 14:1.
- 65. **Wickramasinghe ASD, Kalansuriya P, Attanayake AP.** 2021. Herbal medicines targeting the improved β-cell functions and β-cell regeneration for the management of diabetes mellitus. Evid Based Complement Alternat Med **2021**:2920530.
- 66. Xu X, D'Hoker J, Stangé G, Bonné S, De Leu N, Xiao X, Van de Casteele M, et al. 2008. β Cells can be generated from endogenous progenitors in injured adult mouse pancreas. Cell 132:197–207.
- 67. Yang K, Chan CB. 2018. Epicatechin potentiation of glucosestimulated insulin secretion in INS-1 cells is not dependent on its antioxidant activity. Acta Pharmacol Sin **39**:893–902.
- Yu W, Wang Y, Jiang D, Shang J, Liu M, Efferth T, Teng CB. 2022. A saponin from astragalus promotes pancreatic ductal organoids differentiation into insulin-producing cells. Phytomedicine 102:154190.
- 69. Zaky AS, Kandeil M, Abdel-Gabbar M, Fahmy EM, Almehmadi MM, Ali TM, Ahmed OM. 2022. The antidiabetic effects and modes of action of the balanites aegyptiaca fruit and seed aqueous extracts in NA/STZ-induced diabetic rats. Pharmaceutics 14:263.
- Zebua EA, Silalahi J, Julianti E. 2018. Hypoglicemic activity of gambier (*Uncaria gambir* robx.) drinks in alloxan-induced mice. IOP Conf Ser: Earth Environ Sci 122:012088.
- 71. Zhou J, Kang X, Luo Y, Yuan Y, Wu Y, Wang M, Liu D. 2019. Glibenclamide-induced autophagy inhibits its insulin secretionimproving function in β cells. Int J Endocrinol 2019:1265175.